

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/385418839>

Quadri-Polar Fuzzy Fantastic Ideals in BCI-Algebras: A TOPSIS Framework and Application

Article in *European Journal of Pure and Applied Mathematics* · October 2024

DOI: 10.29020/nybg.ejpam.v17i4.5429

CITATION

1

READS

43

5 authors, including:



M. Balamurugan

Vel Tech Rangarajan Dr. Sagunthala R&D Institute of Science and Technology

25 PUBLICATIONS 188 CITATIONS

[SEE PROFILE](#)



Khalil Hadi Hakami

Jazan University

24 PUBLICATIONS 55 CITATIONS

[SEE PROFILE](#)



Moin Akhtar Ansari

Jazan University

66 PUBLICATIONS 620 CITATIONS

[SEE PROFILE](#)



Anas Al-Masarwah

Ajloun National University

28 PUBLICATIONS 368 CITATIONS

[SEE PROFILE](#)



Quadri-Polar Fuzzy Fantastic Ideals in BCI-Algebras: A TOPSIS Framework and Application

M. Balamurugan¹, Khalil H. Hakami^{2,*}, Moin A. Ansari^{2,*}, Anas Al-Masarwah³,
K. Loganathan⁴

¹ Department of Mathematics, Vel Tech Rangarajan Dr. Sagunthala R&D Institute of Science and Technology, Chennai, Tamil Nadu, India

² Department of Mathematics, College of Science, Jazan University, P.O. Box. 114, Jazan 45142, Kingdom of Saudi Arabia

³ Department of Mathematics, Faculty of Science, Ajloun National University, P.O. Box 43, Ajloun 26810, Jordan

⁴ Department of Mathematics and Statistics, Manipal University Jaipur, Jaipur-303007, India

Abstract. A quadri-polar fuzzy ($q\mathcal{P}\text{-}\mathcal{F}$) set is an extension of a traditional fuzzy set that uses four degrees of membership to represent different aspects of belonging to provide a more detailed framework for handling uncertainty and vagueness. In this paper, we propose the notion of quadri-polar- (ϖ, ϑ) -fuzzy fantastic ideals ($q\mathcal{P}\text{-}(\varpi, \vartheta)\text{-}\mathcal{FFI}(s)$) in BCI-algebras based on $q\mathcal{P}\text{-}\mathcal{F}$ set. Also, the notion of quadri-polar- $(\in_{\tilde{\sigma}}, \in_{\tilde{\sigma}} \vee q_{\tilde{\tau}})$ -fuzzy fantastic ideals ($q\mathcal{P}\text{-}(\in_{\tilde{\sigma}}, \in_{\tilde{\sigma}} \vee q_{\tilde{\tau}})\text{-}\mathcal{FFI}(s)$) is introduced, and the characterizations for an $\in_{\tilde{\sigma}}\text{-}q\mathcal{P}\text{-}\mathcal{F}$ set and $q_{\tilde{\tau}}\text{-}q\mathcal{P}\text{-}\mathcal{F}$ set to be quadri-polar fuzzy ideals ($q\mathcal{P}\text{-}\mathcal{FI}$) in BCI-algebras are established. Furthermore, we present the $q\mathcal{P}\text{-}\mathcal{F}$ TOPSIS technique for multi-criteria Group decision-making (MCGDM), which is a natural extension of the TOPSIS method and used to rank and choose the best alternatives under $q\mathcal{P}\text{-}\mathcal{F}$ positive and negative ideal solutions. Finally, practical examples interpreting the applicability of our proposed $q\mathcal{P}\text{-}\mathcal{F}$ -TOPSIS are solved.

2020 Mathematics Subject Classifications: 03B47, 03E72, 08A72

Key Words and Phrases: BCK/BCI-algebras, q -polar fuzzy fantastic ideal, q -polar- (ω, ϑ) -fuzzy fantastic ideal, q -polar- $(\in_{\tilde{\sigma}}, \in_{\tilde{\sigma}} \vee q_{\tilde{\tau}})$ -fuzzy fantastic ideal

*Corresponding author.

DOI: <https://doi.org/10.29020/nybg.ejpam.v17i4.5429>

Email addresses: drbalamuruganm@veltech.edu.in (M. Balamurugan),
khakami@jazanu.edu.sa (Khalil H. Hakami),
maansari@jazanu.edu.sa (Moin A. Ansari),
anas.almasarwah@anu.edu.jo (Anas Al-Masarwah),
loganathankaruppusamy304@gmail.com (K. Loganathan)

1. Introduction

Axiom systems, developed by Imai et al. [20, 21] and used in propositional calculi, are collections of axioms and inference guidelines used to derive theorems and prove the correctness of logical arguments. Theorem logic and propositional calculus are other names for propositional logic, which deals with the manipulation and analysis of statements using logical operators like OR, AND, and NOT. Various mathematical systems, including propositional logic, can be modeled and analyzed using algebraic structures, particularly Boolean algebra, which is closely related to propositional logic. Iseki [22, 23] introduced the concept of BCK/BCI-algebras. BCI-algebras, also known as BCK-algebras, generalize Boolean algebras and other related algebraic structures. The idea of fantastic ideals in BCI-algebras is a significant algebraic substructure presented and discussed by Saeid [1].

Zadeh [39] proposed the concept of fuzzy (uncertainty) sets, which address ambiguity and vagueness in real world circumstances. A membership function with a range of $[0, 1]$ is used to illustrate an uncertainty structure. Throughout the history of uncertainty set, there are many kinds of uncertainty set extensions, for example bipolar [13] and multipolar [3] uncertainty sets, etc. The bipolar and multipolar uncertainty sets are in fact a generalization of an uncertainty set with a membership degree range $[-1, 1]$ and $[0, 1]^q$, respectively. In [5, 17], the few aspects of the bipolar fuzzy concept are applied to algebraic structures. The $q\mathcal{P}\text{-}\mathcal{FS}$ has an extensive range of implementations to address ambiguity and vagueness in real world issues related to the quadri-polar data, quadri-index and quadri-attributes information. Researchers in a lot of different areas are very interested in the multi-polar uncertainty set theory. These areas include Lie algebras [4], ordered semihypergroups [30], subgroups [16] and BCK/BCI-algebras [7, 36].

Rosenfeld [38] introduced fuzzy groups, while Bhakat et al [12] developed a specific type denoted as $(\in, \in \vee q)$, based on point fuzzy sets within group theory. Jun [27, 28] and Muhiuddin et al. [35] extended this concept to (α, β) -fuzzy subalgebra. Ibrara et al. [19], Dudek et al. [14], and Narayanan et al. [37] furthered this idea with extensions to semigroups, hemirings, and near-rings, respectively. Al-Masarwah et al. [6, 8] explored (α, β) type subalgebras using m-F points within BCK-algebras. Ma et al. [33] introduced $(\in_\gamma, \in_\gamma \vee q_\delta)$ -fuzzy ideals, while Jana et al. [24] proposed $(\in_\gamma, \in_\gamma \vee q_\delta)$ fuzzy soft BCI-algebras. Zulfiqar et al. [42, 43] introduced the idea of $(\in_\gamma, \in_\gamma \vee q_\delta)$ -fuzzy subcommutative ideals and fuzzy fantastic ideals in BCI/BCH-algebras. Zhan [41] contributed with $(\in_\gamma, \in_\gamma \vee q_\delta)$ -fuzzy soft Γ -hyper ideals. Abuhijleh et al. [2] introduced the complex fuzzy groups. Fallath et al. [15] introduced cosets and normals of (γ, δ) -fuzzy HX-subgroups. Balamurugan et al. [10, 18] introduced anti-intuitionistic fuzzy soft ideals in BCK/BCI/BG-algebras. Balamurugan et al. [11, 34] introduced tripolar picture fuzzy ideals and bipolar intuitionistic fuzzy soft ideals in BCK/BCI-algebras. Moin et al. [9] introduced and studied a graph associated to UP-algebras. Fuzzy bi-ideals in ternary semirings are studied and explored by Kavikumar [29].

In this work, we combine $q\mathcal{P}\text{-}\mathcal{F}$ sets with BCI-algebras to extend fuzzy set theory and provide new approaches for studying quadri-polar fuzzy BCI-algebras. We introduce a new class of generalized $q\mathcal{P}\text{-}(\varpi, \vartheta)\text{-}\mathcal{FFI}$. The properties of $q\mathcal{P}\text{-}(\varpi, \vartheta)\text{-}\mathcal{FFI}(s)$ are highlighted.

We then discuss $q\mathcal{P}-(\in_{\tilde{\sigma}}, \in_{\tilde{\sigma}} \vee q_{\tilde{\tau}})\mathcal{FFI}(s)$ and explore their properties. Characterization theorems for $q\mathcal{P}-(\in_{\tilde{\sigma}}, \in_{\tilde{\sigma}} \vee q_{\tilde{\tau}})\mathcal{FFI}(s)$ are also established. Finally, we present a $q\mathcal{PF}$ TOPSIS methodology, discuss potential applications, compare it with existing TOPSIS methods, and propose future directions.

To explain the novelty of this structure, some contributions by several researchers towards $q\mathcal{P}\text{-}\mathcal{FFI}(s)$, $q\mathcal{P}-(\varpi, \vartheta)\text{-}\mathcal{FFI}(s)$ and $q\mathcal{P}-(\in_{\tilde{\sigma}}, \in_{\tilde{\sigma}} \vee q_{\tilde{\tau}})\text{-}\mathcal{FFI}(s)$ in BCI-algebras are presented in Table 1.

Table 1: Contributions of several researchers toward ceratin generalizations of $q\mathcal{P}\text{-}\mathcal{FFI}(s)$.

Authors	Year	Contributions
Rosenfeld [38]	1971	Creation of fuzzy subgroups.
Xi [40]	1991	Creation of fuzzy ideals.
Bhakat and Das [12]	1996	Certain extensions of fuzzy subgroups.
Jun [26]	2004	Creation of (α, β) -fuzzy ideals.
Lee [32]	2009	Creation of bipolar fuzzy ideals.
Jana et al. [25]	2017	Extensions of bipolar fuzzy ideals.
Al-Masarwah and Ahmad [6–8]	2018	Creation of multi $\mathcal{P}\text{-}\mathcal{FI}(s)$.
Alqahtani et al.	Present	Creation of $q\mathcal{P}-(\in_{\tilde{\sigma}}, \in_{\tilde{\sigma}} \vee q_{\tilde{\tau}})\text{-}\mathcal{FFI}(s)$.

2. Preliminaries

BCI-algebras are types of algebraic structures used in the study of non-classical logics, particularly in the context of certain types of implication algebras. These algebras generalize certain aspects of set theory, logic and have applications in some areas, such as theoretical computer science and mathematical logic.

A BCI-algebra is a structure $(\tilde{\mathfrak{N}}; \tilde{\varnothing}, 0)$ consisting of a non-void set $\tilde{\mathfrak{N}}$, a binary operation $\tilde{\varnothing}$ on $\tilde{\mathfrak{N}}$, and a constant $0 \in \tilde{\mathfrak{N}}$, satisfying the following axioms: $\forall \zeta, \dot{\varrho}, \dot{\kappa} \in \tilde{\mathfrak{N}}$

$$(I_1) ((\zeta \tilde{\varnothing} \dot{\varrho}) \tilde{\varnothing} (\zeta \tilde{\varnothing} \dot{\kappa})) \tilde{\varnothing} (\dot{\kappa} \tilde{\varnothing} \dot{\varrho}) = 0,$$

$$(I_2) (\zeta \tilde{\varnothing} (\zeta \tilde{\varnothing} \dot{\varrho})) \tilde{\varnothing} \dot{\varrho} = 0,$$

$$(I_3) \zeta \tilde{\varnothing} \zeta = 0,$$

$$(I_4) \zeta \tilde{\varnothing} \dot{\varrho} = 0, \dot{\varrho} \tilde{\varnothing} \zeta = 0 \Rightarrow \zeta = \dot{\varrho}.$$

A subset \mathcal{I} of $\tilde{\mathfrak{N}}$ is referred to an *ideal* of $\tilde{\mathfrak{N}}$ (see [22, 23]) if it meets:

$$0 \in \mathcal{I} \text{ and } (\forall \zeta, \dot{\varrho} \in \mathcal{I}) (\zeta \tilde{\varnothing} \dot{\varrho} \in \mathcal{I}, \dot{\varrho} \in \mathcal{I} \Rightarrow \zeta \in \mathcal{I}). \quad (1)$$

A subset \mathcal{I} of $\tilde{\mathfrak{N}}$ is referred to a *fantastic ideal* of $\tilde{\mathfrak{N}}$ (see [1]) if it meets:

$$0 \in \mathcal{I} \text{ and } (\forall \zeta, \dot{\varrho}, \dot{\kappa} \in \mathcal{I}) ((\zeta \tilde{\varnothing} \dot{\varrho}) \tilde{\varnothing} \dot{\kappa} \in \mathcal{I}, \dot{\kappa} \in \mathcal{I} \Rightarrow \zeta \tilde{\varnothing} (\dot{\varrho} \tilde{\varnothing} (\dot{\varrho} \tilde{\varnothing} \zeta)) \in \mathcal{I}). \quad (2)$$

Definition 1. [31] A mapping $\tilde{A} : \tilde{\mathfrak{N}} \rightarrow [0, 1]$ is a fuzzy set \mathcal{FS} for the set $\tilde{\mathfrak{N}}$.

A $\mathcal{FS} \tilde{A}$ of \tilde{N} is a \mathcal{FI} of \tilde{N} if it meets:

$$(\forall \zeta, \varrho \in \tilde{N}, \tilde{A}(0) \geq \tilde{A}(\zeta) \text{ and } \tilde{A}(\zeta) \geq \tilde{A}(\zeta \oslash \varrho) \wedge \tilde{A}(\varrho)). \quad (3)$$

A $\mathcal{FS} \tilde{A}$ of \tilde{N} is a \mathcal{FFI} of \tilde{N} if it meets:

$$(\forall \zeta, \varrho \in \tilde{N}, \tilde{A}(0) \geq \tilde{A}(\zeta) \text{ and } \tilde{A}(\zeta \oslash (\varrho \oslash (\varrho \oslash \zeta))) \geq \tilde{A}((\zeta \oslash \varrho) \oslash \kappa) \wedge \tilde{A}(\kappa)). \quad (4)$$

3. Quadri-Polar Fuzzy Fantastic Ideals

Definition 2. A mapping $\tilde{\theta} : \tilde{N} \rightarrow [0, 1]^4$ is a $q\mathcal{P}\text{-}\mathcal{F}$ for the set \tilde{N} , where for any $\zeta \in \tilde{N}$,

$$\tilde{\theta}(\zeta) = (\tilde{\theta}^1(\zeta), \tilde{\theta}^2(\zeta), \tilde{\theta}^3(\zeta), \tilde{\theta}^4(\zeta)) \text{ and } \tilde{\theta}^q(\zeta) \in [0, 1],$$

for $q = 1, 2, 3, 4$.

Definition 3. A $q\mathcal{P}\text{-}\mathcal{F}$ set $\tilde{\theta}$ of \tilde{N} is a $q\mathcal{P}\text{-}\mathcal{FFI}$ if, $\forall \zeta, \varrho, \kappa \in \tilde{N}$ and $q = 1, 2, 3, 4$,

$$\tilde{\theta}(0) \geq \tilde{\theta}(\zeta) \text{ and } \tilde{\theta}(\zeta \oslash (\varrho \oslash (\varrho \oslash \zeta))) \geq \tilde{\theta}((\zeta \oslash \varrho) \oslash \kappa) \wedge \tilde{\theta}(\kappa).$$

That is,

$$\tilde{\theta}^q(0) \geq \tilde{\theta}^q(\zeta) \text{ and } \tilde{\theta}^q(\zeta \oslash (\varrho \oslash (\varrho \oslash \zeta))) \geq \tilde{\theta}^q((\zeta \oslash \varrho) \oslash \kappa) \wedge \tilde{\theta}^q(\kappa).$$

Example 1. Consider $\tilde{N} = \{0, \zeta, \varrho, \kappa\}$ with the binary operation \oslash defined by Table 2:

Table 2. Cayley table representing by “ \oslash ”

\oslash	0	ζ	ϱ	κ
0	0	0	0	0
ζ	ζ	0	0	ζ
ϱ	ϱ	ζ	0	ϱ
κ	κ	κ	κ	0

Thus, $(\tilde{N}; \oslash, 0)$ forms a BCI-algebra.

Consider a $q\mathcal{P}\text{-}\mathcal{F}$ set $\tilde{\theta}$ defined on \tilde{N} as follows:

$$\tilde{\theta}(\zeta) = \left\{ \begin{array}{l} \langle 0, (.58, .65, .75, .54) \rangle, \\ \langle \zeta, (.48, .21, .45, .30) \rangle, \\ \langle \varrho, (.28, .52, .54, .30) \rangle, \\ \langle \kappa, (.28, .41, .36, .54) \rangle. \end{array} \right\}$$

Thus, $\tilde{\theta}$ is a $q\mathcal{P}\text{-}\mathcal{FFI}$ of \tilde{N} .

Theorem 1. A $q\mathcal{P}\text{-}\mathcal{F}$ set $\tilde{\theta}$ is a $q\mathcal{P}\text{-}\mathcal{FFI}$ of $\tilde{N} \Leftrightarrow$ for any $\tilde{\rho} \in (0, 1]^4$, the $\tilde{\rho}$ -cut subset $\tilde{\theta}_{\tilde{\rho}} = \{\zeta \in \tilde{N} \mid \tilde{\theta}(\zeta) \geq \tilde{\rho}\}$ is a fantastic ideal of \tilde{N} .

Proof. Let $\tilde{\theta}$ be a $q\mathcal{P}\text{-}\mathcal{FFI}$ of \tilde{N} and $\tilde{\rho} \in (0, 1]^4$ be such that $\tilde{\theta}_{\tilde{\rho}} = \{\zeta \in \tilde{N} \mid \tilde{\theta}(\zeta) \geq \tilde{\rho}\}$. Let $\zeta, \varrho, \kappa \in \tilde{\theta}_{\tilde{\rho}}$. Then, $\tilde{\theta}((\zeta \oslash \varrho) \oslash \kappa) \geq \tilde{\rho}$ and $\tilde{\theta}(\kappa) \geq \tilde{\rho}$. It follows from Definition 3.2 that,

$$\tilde{\mathfrak{d}}(\zeta \mathbin{\dot{\vee}} (\varrho \mathbin{\dot{\vee}} (\varrho \mathbin{\dot{\vee}} \zeta))) \geq \tilde{\mathfrak{d}}((\zeta \mathbin{\dot{\vee}} \varrho) \mathbin{\dot{\vee}} \kappa) \wedge \tilde{\mathfrak{d}}(\kappa) = \tilde{\rho} \wedge \tilde{\rho} = \tilde{\rho}.$$

Therefore, $\zeta \mathbin{\dot{\vee}} (\varrho \mathbin{\dot{\vee}} (\varrho \mathbin{\dot{\vee}} \zeta)) \in \tilde{\mathfrak{d}}_{\tilde{\rho}}$. Hence, $\tilde{\mathfrak{d}}_{\tilde{\rho}}$ is a fantastic ideal of $\tilde{\mathfrak{N}}$.

Conversely, assume $\tilde{\mathfrak{d}}_{\tilde{\rho}}$ is a fantastic ideal of $\tilde{\mathfrak{N}}$. Suppose that $\tilde{\mathfrak{d}}(\zeta \mathbin{\dot{\vee}} (\varrho \mathbin{\dot{\vee}} (\varrho \mathbin{\dot{\vee}} \zeta))) < \tilde{\mathfrak{d}}((\zeta \mathbin{\dot{\vee}} \varrho) \mathbin{\dot{\vee}} \kappa) \wedge \tilde{\mathfrak{d}}(\kappa)$. Then $\tilde{\mathfrak{d}}(\zeta \mathbin{\dot{\vee}} (\varrho \mathbin{\dot{\vee}} (\varrho \mathbin{\dot{\vee}} \zeta))) < \tilde{\rho} \leq \tilde{\mathfrak{d}}((\zeta \mathbin{\dot{\vee}} \varrho) \mathbin{\dot{\vee}} \kappa) \wedge \tilde{\mathfrak{d}}(\kappa)$. But $\tilde{\mathfrak{d}}((\zeta \mathbin{\dot{\vee}} \varrho) \mathbin{\dot{\vee}} \kappa) \geq \tilde{\rho}$ and $\tilde{\mathfrak{d}}(\kappa) \geq \tilde{\rho}$. So, $\zeta \mathbin{\dot{\vee}} (\varrho \mathbin{\dot{\vee}} (\varrho \mathbin{\dot{\vee}} \zeta)) \notin \tilde{\mathfrak{d}}_{\tilde{\rho}}$, a contradiction. Therefore,

$$\tilde{\mathfrak{d}}(\zeta \mathbin{\dot{\vee}} (\varrho \mathbin{\dot{\vee}} (\varrho \mathbin{\dot{\vee}} \zeta))) \geq \tilde{\mathfrak{d}}((\zeta \mathbin{\dot{\vee}} \varrho) \mathbin{\dot{\vee}} \kappa) \wedge \tilde{\mathfrak{d}}(\kappa).$$

Hence, $\tilde{\mathfrak{d}}_{\tilde{\rho}}$ is a \mathcal{FFI} of $\tilde{\mathfrak{N}}$.

Consider a $q\mathcal{P}$ - \mathcal{F} set $\tilde{\mathfrak{d}}$ defined on $\tilde{\mathfrak{N}}$, where

$$\tilde{\mathfrak{d}}(\zeta) = \begin{cases} \tilde{\rho} \in (0, 1]^q, & \text{if } \zeta \in \tilde{\mathfrak{N}} \\ 0, & \text{if } \zeta \notin \tilde{\mathfrak{N}}, \end{cases}$$

then $\tilde{\mathfrak{d}}(\zeta)$ is a $q\mathcal{P}$ - \mathcal{F} point with support $\tilde{\mathfrak{N}}$ and the value $\tilde{\rho}$, and it is symbolized by $\zeta_{\tilde{\rho}}$.

Theorem 2. Every fantastic ideal of $\tilde{\mathfrak{N}}$ is a $q\mathcal{P}$ - \mathcal{FFI} of $\tilde{\mathfrak{N}}$.

Proof. Suppose $\tilde{\mathfrak{d}}_{\tilde{\rho}}$ is a fantastic ideal of $\tilde{\mathfrak{N}}$ and let $\tilde{\mathfrak{d}}$ be an $q\mathcal{P}$ - \mathcal{FS} in $\tilde{\mathfrak{N}}$ defined by

$$\tilde{\mathfrak{d}}(\zeta) = \begin{cases} \tilde{\rho} \in (0, 1]^q, & \text{if } \zeta \in \tilde{\mathfrak{N}} \\ 0, & \text{if } \zeta \notin \tilde{\mathfrak{N}} \end{cases}$$

Let $\zeta, \varrho \in \tilde{\mathfrak{N}}$. To verify that $\tilde{\mathfrak{d}}$ is a $q\mathcal{P}$ - \mathcal{FFI} of $\tilde{\mathfrak{N}}$.

Case 1: If $(\zeta \mathbin{\dot{\vee}} \varrho) \mathbin{\dot{\vee}} \kappa \in \tilde{\mathfrak{d}}$ and $\kappa \in \tilde{\mathfrak{d}}$, then $(\zeta \mathbin{\dot{\vee}} (\varrho \mathbin{\dot{\vee}} (\varrho \mathbin{\dot{\vee}} \zeta))) \in \tilde{\mathfrak{d}}$. Thus $\tilde{\mathfrak{d}}((\zeta \mathbin{\dot{\vee}} \varrho) \mathbin{\dot{\vee}} \kappa) = \tilde{\mathfrak{d}}(\kappa) = \tilde{\rho}$. Hence by Definition 3.2, we have

$$\tilde{\mathfrak{d}}(\zeta \mathbin{\dot{\vee}} (\varrho \mathbin{\dot{\vee}} (\varrho \mathbin{\dot{\vee}} \zeta))) \geq \tilde{\mathfrak{d}}((\zeta \mathbin{\dot{\vee}} \varrho) \mathbin{\dot{\vee}} \kappa) \wedge \tilde{\mathfrak{d}}(\kappa) = \tilde{\rho} \wedge \tilde{\rho} = \tilde{\rho}.$$

Case 2: If $(\zeta \mathbin{\dot{\vee}} \varrho) \mathbin{\dot{\vee}} \kappa \notin \tilde{\mathfrak{d}}$ and $\kappa \notin \tilde{\mathfrak{d}}$, then $(\zeta \mathbin{\dot{\vee}} (\varrho \mathbin{\dot{\vee}} (\varrho \mathbin{\dot{\vee}} \zeta))) \notin \tilde{\mathfrak{d}}$. Thus, $\tilde{\mathfrak{d}}((\zeta \mathbin{\dot{\vee}} \varrho) \mathbin{\dot{\vee}} \kappa) = \tilde{\mathfrak{d}}(\kappa) = 0$. Hence by Definition 3.2, we have

$$\tilde{\mathfrak{d}}(\zeta \mathbin{\dot{\vee}} (\varrho \mathbin{\dot{\vee}} (\varrho \mathbin{\dot{\vee}} \zeta))) \geq \tilde{\mathfrak{d}}((\zeta \mathbin{\dot{\vee}} \varrho) \mathbin{\dot{\vee}} \kappa) \wedge \tilde{\mathfrak{d}}(\kappa) = 0 \wedge 0 = 0.$$

Case 3: If either $(\zeta \mathbin{\dot{\vee}} \varrho) \mathbin{\dot{\vee}} \kappa \in \tilde{\mathfrak{d}}$ or $\kappa \in \tilde{\mathfrak{d}}$, then either $\tilde{\mathfrak{d}}((\zeta \mathbin{\dot{\vee}} \varrho) \mathbin{\dot{\vee}} \kappa) = \tilde{\rho}$ or $\tilde{\mathfrak{d}}(\kappa) = \tilde{\rho}$. So,

$$\tilde{\mathfrak{d}}(\zeta \mathbin{\dot{\vee}} (\varrho \mathbin{\dot{\vee}} (\varrho \mathbin{\dot{\vee}} \zeta))) \geq \tilde{\mathfrak{d}}((\zeta \mathbin{\dot{\vee}} \varrho) \mathbin{\dot{\vee}} \kappa) \wedge \tilde{\mathfrak{d}}(\kappa).$$

Hence, $\tilde{\mathfrak{d}}$ is a $q\mathcal{P}$ - \mathcal{FFI} of $\tilde{\mathfrak{N}}$.

4. Quadri-Polar (ϖ, ϑ) -Fuzzy Fantastic Ideals

In this section, we introduce the concept of a $q\mathcal{P}-(\varpi, \vartheta)\mathcal{FFI}(s)$ in BCI-algebras and explore various properties associated with it. Here, we use ϖ and ϑ to represent symbols such as $\in_{\tilde{\sigma}}, \in_{\tilde{\sigma}} \vee q_{\tilde{\tau}}, q_{\tilde{\tau}}$ or $\in_{\tilde{\sigma}} \wedge q_{\tilde{\tau}}$, unless specified otherwise.

Consider a $q\mathcal{P}\text{-}\mathcal{F}$ point denoted as $\zeta_{\tilde{\rho}}$ and a $q\mathcal{P}\text{-}\mathcal{F}$ set $\tilde{\mathfrak{O}}$ defined on $\tilde{\aleph}$. Then

- (1) $\zeta_{\tilde{\rho}} \in_{\tilde{\sigma}} \tilde{\mathfrak{O}}$ if $\tilde{\mathfrak{O}}(\zeta) \geq \tilde{\rho} > \tilde{\sigma}$.
 - (2) $\zeta_{\tilde{\rho}} q_{\tilde{\tau}} \tilde{\mathfrak{O}}$ if $\tilde{\mathfrak{O}}(\zeta) + \tilde{\rho} > 2\tilde{\tau}$.
 - (3) $\zeta_{\tilde{\rho}} \in_{\tilde{\sigma}} \vee q_{\tilde{\tau}} \tilde{\mathfrak{O}}$ if $\zeta_{\tilde{\rho}} \in_{\tilde{\sigma}} \tilde{\mathfrak{O}}$ or $\zeta_{\tilde{\rho}} q_{\tilde{\tau}} \tilde{\mathfrak{O}}$.
 - (4) $\zeta_{\tilde{\rho}} \in_{\tilde{\sigma}} \wedge q_{\tilde{\tau}} \tilde{\mathfrak{O}}$ if $\zeta_{\tilde{\rho}} \in_{\tilde{\sigma}} \tilde{\mathfrak{O}}$ and $\zeta_{\tilde{\rho}} q_{\tilde{\tau}} \tilde{\mathfrak{O}}$.
 - (5) $\zeta_{\tilde{\rho}} \varpi \tilde{\mathfrak{O}}$ does not hold for $\varpi = \{\in_{\tilde{\sigma}}, q_{\tilde{\tau}}, \in_{\tilde{\sigma}} \vee q_{\tilde{\tau}}, \in_{\tilde{\sigma}} \wedge q_{\tilde{\tau}}\}, \forall \tilde{\sigma}, \tilde{\tau} \in [0, 1]^4$, where $\tilde{\sigma} = (\tilde{\sigma}_1, \tilde{\sigma}_2, \tilde{\sigma}_3, \tilde{\sigma}_4) < \tilde{\tau} = (\tilde{\tau}_1, \tilde{\tau}_2, \tilde{\tau}_3, \tilde{\tau}_4)$.
- A $q\mathcal{P}\text{-}\mathcal{F}$ point $\zeta_{\tilde{\rho}} \in \tilde{\mathfrak{O}}$ if $\tilde{\mathfrak{O}}(\zeta) \geq \tilde{\rho}$. That is $\tilde{\mathfrak{O}}^q(\zeta) \geq \tilde{\rho}^q, \forall q = 1, 2, 3, 4$. Also, $\zeta_{\tilde{\rho}} q \tilde{\mathfrak{O}}$ if $\tilde{\mathfrak{O}}(\zeta) + \tilde{\rho} > \hat{1}$. That is, $\tilde{\mathfrak{O}}^q(\zeta) + \tilde{\rho}^q > \hat{1}, \forall q = 1, 2, 3, 4$.

By $\zeta_{\tilde{\rho}} \in \vee q \tilde{\mathfrak{O}}$ (resp., $\zeta_{\tilde{\rho}} \in \wedge q \tilde{\mathfrak{O}}$) $\Rightarrow \zeta_{\tilde{\rho}} \in \tilde{\mathfrak{O}}$ or $\zeta_{\tilde{\rho}} q \tilde{\mathfrak{O}}$ (resp., $\zeta_{\tilde{\rho}} \in \tilde{\mathfrak{O}}$ and $\zeta_{\tilde{\rho}} q \tilde{\mathfrak{O}}$). If $\phi \neq \tilde{C} \subseteq \tilde{\aleph}$, then the quadri-polar characteristic fuzzy set ($q\mathcal{P}\text{-}\mathcal{CF}$) of \tilde{C} , say $\hat{\chi}_{\tilde{C}}$, where

$$\hat{\chi}_{\tilde{C}} = \begin{cases} \hat{1} = (1, 1, 1, 1), & \text{if } \zeta \in \tilde{C} \\ \tilde{0} = (0, 0, 0, 0), & \text{if } \zeta \notin \tilde{C} \end{cases}$$

Clearly, a $q\mathcal{P}\text{-}\mathcal{CF}$ is a $q\mathcal{P}\text{-}\mathcal{F}$ subset of $\tilde{\aleph}$.

Definition 4. A $q\mathcal{P}\text{-}\mathcal{F}$ set $\tilde{\mathfrak{O}}$ is a $q\mathcal{P}-(\varpi, \vartheta)\text{-}\mathcal{FFI}$ of $\tilde{\aleph}$, if

$$((\zeta \check{\vee} \varrho) \check{\vee} \kappa)_{\tilde{\rho}} \varpi \tilde{\mathfrak{O}}, \kappa_{\tilde{\eta}} \varpi \tilde{\mathfrak{O}} \Rightarrow (\zeta \check{\vee} (\varrho \check{\vee} (\varrho \check{\vee} \zeta)))_{\tilde{\rho} \wedge \tilde{\eta}} \vartheta \tilde{\mathfrak{O}},$$

where $\varpi \neq \in_{\tilde{\sigma}} \wedge q_{\tilde{\tau}}, \forall \tilde{\sigma} < \tilde{\rho}, \tilde{\eta} \leq \hat{1}$ and $((\zeta \check{\vee} \varrho) \check{\vee} \kappa), \kappa \in \tilde{\aleph}$.

Consider a $q\mathcal{P}\text{-}\mathcal{F}$ set $\tilde{\mathfrak{O}}$ defined on $\tilde{\aleph}$ such that $\tilde{\mathfrak{O}}(\zeta) \leq \tilde{\tau}, \forall \zeta \in \tilde{\aleph}$. Let $\zeta \in \tilde{\aleph}$ and $\tilde{\sigma} < \tilde{\rho} \leq \hat{1}$ be such that $\zeta_{\tilde{\rho}} \in \wedge q_{\tilde{\tau}} \tilde{\mathfrak{O}}$. Then, $\tilde{\mathfrak{O}}(\zeta) \geq \tilde{\rho} > \tilde{\sigma}$ and $\tilde{\mathfrak{O}}(\zeta) + \tilde{\rho} > 2\tilde{\tau}$. Thus,

$$2\tilde{\tau} < \tilde{\mathfrak{O}}(\zeta) + \tilde{\rho} \leq \tilde{\mathfrak{O}}(\zeta) + \tilde{\mathfrak{O}}(\zeta) = 2\tilde{\mathfrak{O}}(\zeta) \Rightarrow \tilde{\mathfrak{O}}(\zeta) > \tilde{\tau}.$$

Hence, $\{\zeta_{\tilde{\rho}} \mid \zeta_{\tilde{\rho}} \in_{\tilde{\sigma}} \wedge q_{\tilde{\tau}} \tilde{\mathfrak{O}}\} = \emptyset$. Therefore, we exclude the case $\varpi = \in_{\tilde{\sigma}} \wedge q_{\tilde{\tau}}$ in Definition 4.1 is neglected.

Theorem 3. Let $\tilde{\mathfrak{O}}$ be a $q\mathcal{P}-(\varpi, \vartheta)\text{-}\mathcal{FFI}$ and $\tilde{\sigma} + \hat{1} = 2\tilde{\tau}$ of $\tilde{\aleph}$. Then, the set

$$\tilde{\mathfrak{O}}_{\tilde{\sigma}} = \{\zeta \in \tilde{\aleph} \mid \tilde{\mathfrak{O}}(\zeta) > \tilde{\sigma}\}$$

is a \mathcal{FI} of $\tilde{\aleph}$.

Proof. Let $\zeta, \varrho, \kappa \in \tilde{\aleph}$ be such that $\zeta, \varrho, \kappa \in \tilde{\mathfrak{O}}_{\tilde{\sigma}}$. Then, $\tilde{\mathfrak{O}}((\zeta \check{\vee} \varrho) \check{\vee} \kappa) > \tilde{\sigma}$ and $\tilde{\mathfrak{O}}(\kappa) > \tilde{\sigma}$. Assume $\tilde{\mathfrak{O}}(\zeta \check{\vee} (\varrho \check{\vee} (\varrho \check{\vee} \zeta))) \leq \tilde{\sigma}$. If $\varpi \in \{\in_{\tilde{\sigma}}, \in_{\tilde{\sigma}} \vee q_{\tilde{\tau}}\}$, then

$$((\dot{\varsigma} \check{\vee} \dot{\varrho}) \check{\vee} \dot{\kappa})_{\check{\partial}(\dot{\varsigma})} \varpi \check{\partial} \text{ and } \dot{\kappa}_{\check{\partial}(\dot{\varrho})} \varpi \check{\partial}.$$

But

$$\check{\partial}(\dot{\varsigma} \check{\vee} (\dot{\varrho} \check{\vee} (\dot{\varrho} \check{\vee} \dot{\varsigma}))) \leq \tilde{\sigma} < \check{\partial}((\dot{\varsigma} \check{\vee} \dot{\varrho}) \check{\vee} \dot{\kappa}) \wedge \check{\partial}(\dot{\kappa})$$

and

$$\check{\partial}(\dot{\varsigma} \check{\vee} (\dot{\varrho} \check{\vee} (\dot{\varrho} \check{\vee} \dot{\varsigma}))) + \check{\partial}((\dot{\varsigma} \check{\vee} \dot{\varrho}) \check{\vee} \dot{\kappa}) \wedge \check{\partial}(\dot{\kappa}) \leq \tilde{\sigma} + \hat{1} = 2\tilde{\tau}.$$

So,

$$(\dot{\varsigma} \check{\vee} (\dot{\varrho} \check{\vee} (\dot{\varrho} \check{\vee} \dot{\varsigma})))_{\check{\partial}(\dot{\varsigma}) \wedge \check{\partial}(\dot{\eta})} \vartheta \check{\partial}, \forall \vartheta \in \{\in_{\tilde{\sigma}}, q_{\tilde{\tau}}, \in_{\tilde{\sigma}} \vee q_{\tilde{\tau}}, \in_{\tilde{\sigma}} \wedge q_{\tilde{\tau}}\}, \text{ a contradiction.}$$

Hence, $\check{\partial}(\dot{\varsigma} \check{\vee} (\dot{\varrho} \check{\vee} (\dot{\varrho} \check{\vee} \dot{\varsigma}))) > \tilde{\sigma} \Rightarrow \dot{\varsigma} \check{\vee} (\dot{\varrho} \check{\vee} (\dot{\varrho} \check{\vee} \dot{\varsigma})) \in \check{\partial}_{\tilde{\sigma}}$.

Also,

$$\check{\partial}(\dot{\varsigma} \check{\vee} (\dot{\varrho} \check{\vee} (\dot{\varrho} \check{\vee} \dot{\varsigma}))) + \hat{1} > \tilde{\sigma} + \hat{1} = 2\tilde{\tau} \Rightarrow (\dot{\varsigma} \check{\vee} (\dot{\varrho} \check{\vee} (\dot{\varrho} \check{\vee} \dot{\varsigma})))_1 q_{\tilde{\tau}} \check{\partial}.$$

But

$$\check{\partial}(\dot{\varsigma} \check{\vee} (\dot{\varrho} \check{\vee} (\dot{\varrho} \check{\vee} \dot{\varsigma}))) \leq \tilde{\sigma} \Rightarrow (\dot{\varsigma} \check{\vee} (\dot{\varrho} \check{\vee} (\dot{\varrho} \check{\vee} \dot{\varsigma})))_1 \overline{\in_{\tilde{\sigma}}} \check{\partial}$$

and

$$\check{\partial}(\dot{\varsigma} \check{\vee} (\dot{\varrho} \check{\vee} (\dot{\varrho} \check{\vee} \dot{\varsigma}))) + \hat{1} \leq \tilde{\sigma} + \hat{1} = 2\tilde{\tau} \Rightarrow (\dot{\varsigma} \check{\vee} (\dot{\varrho} \check{\vee} (\dot{\varrho} \check{\vee} \dot{\varsigma})))_1 \overline{q_{\tilde{\tau}}} \check{\partial}, \text{ a contradiction.}$$

Thus,

$$\check{\partial}(\dot{\varsigma} \check{\vee} (\dot{\varrho} \check{\vee} (\dot{\varrho} \check{\vee} \dot{\varsigma}))) > \tilde{\sigma} \Rightarrow \dot{\varsigma} \check{\vee} (\dot{\varrho} \check{\vee} (\dot{\varrho} \check{\vee} \dot{\varsigma})) \in \check{\partial}_{\tilde{\sigma}}.$$

Therefore, $\check{\partial}_{\tilde{\sigma}}$ is a \mathcal{FI} of $\tilde{\aleph}$.

Theorem 4. Let $\phi \neq \tilde{C} \subseteq \tilde{\aleph}$ and $\tilde{\sigma} + \hat{1} = 2\tilde{\tau}$. Then \tilde{C} is a \mathcal{FI} of $\tilde{\aleph}$ if and only if the $q\mathcal{P}$ - \mathcal{F} subset $\check{\partial}$ of $\tilde{\aleph}$, which is defined as follows:

- (1) $\check{\partial}(\dot{\varsigma}) \geq \tilde{\tau}, \forall \dot{\varsigma} \in \tilde{C}$,
- (2) $\check{\partial}(\dot{\varsigma}) \leq \tilde{\sigma}, \forall \dot{\varsigma} \notin \tilde{C}$ is a $q\mathcal{P}$ -($\in_{\tilde{\sigma}}, \in_{\tilde{\sigma}} \vee q_{\tilde{\tau}}$)- \mathcal{FFI} of $\tilde{\aleph}$.

Proof. Let \tilde{C} be a \mathcal{FI} of $\tilde{\aleph}$, $\dot{\varsigma}, \dot{\varrho}, \dot{\kappa} \in \tilde{\aleph}$ and let $\tilde{\sigma} < \tilde{\rho}, \tilde{\eta} \leq \hat{1}$ be such that

$$((\dot{\varsigma} \check{\vee} \dot{\varrho}) \check{\vee} \dot{\kappa})_{\tilde{\rho}} \in_{\tilde{\sigma}} \vee q_{\tilde{\tau}} \check{\partial} \text{ and } \dot{\kappa}_{\tilde{\eta}} \in_{\tilde{\sigma}} \check{\partial}.$$

Then

$$\check{\partial}(((\dot{\varsigma} \check{\vee} \dot{\varrho}) \check{\vee} \dot{\kappa})) \geq \tilde{\rho} > \tilde{\sigma} \text{ and } \check{\partial}(\dot{\kappa}) \geq \tilde{\eta} > \tilde{\sigma}.$$

Thus,

$$\dot{\varsigma} \check{\vee} (\dot{\varrho} \check{\vee} (\dot{\varrho} \check{\vee} \dot{\varsigma})) \in \tilde{C} \Rightarrow \check{\partial}(\dot{\varsigma} \check{\vee} (\dot{\varrho} \check{\vee} (\dot{\varrho} \check{\vee} \dot{\varsigma}))) \geq \tilde{\tau}.$$

If $\tilde{\rho} \wedge \tilde{\eta} \leq \tilde{\tau}$, then

$$\tilde{\mathfrak{O}}(\zeta \mathbin{\dot{\vee}} (\dot{\rho} \mathbin{\dot{\vee}} (\dot{\rho} \mathbin{\dot{\vee}} \zeta))) \geq \tilde{\tau} \geq \tilde{\rho} \wedge \tilde{\eta} > \tilde{\sigma} \Rightarrow (\zeta \mathbin{\dot{\vee}} (\dot{\rho} \mathbin{\dot{\vee}} (\dot{\rho} \mathbin{\dot{\vee}} \zeta)))_{\tilde{\rho} \wedge \tilde{\eta}} \in_{\tilde{\sigma}} \tilde{\mathfrak{O}}.$$

If $\tilde{\rho} \wedge \tilde{\eta} > \tilde{\tau}$, then

$$\tilde{\mathfrak{O}}(\zeta \mathbin{\dot{\vee}} (\dot{\rho} \mathbin{\dot{\vee}} (\dot{\rho} \mathbin{\dot{\vee}} \zeta))) + \tilde{\rho} \wedge \tilde{\eta} > \tilde{\tau} + \tilde{\tau} = 2\tilde{\tau} \Rightarrow (\zeta \mathbin{\dot{\vee}} (\dot{\rho} \mathbin{\dot{\vee}} (\dot{\rho} \mathbin{\dot{\vee}} \zeta)))_{\tilde{\rho} \wedge \tilde{\eta}} q_{\tilde{\tau}} \tilde{\mathfrak{O}}.$$

Thus,

$$(\zeta \mathbin{\dot{\vee}} (\dot{\rho} \mathbin{\dot{\vee}} (\dot{\rho} \mathbin{\dot{\vee}} \zeta)))_{\tilde{\rho} \wedge \tilde{\eta}} \in_{\tilde{\sigma}} \vee q_{\tilde{\tau}} \tilde{\mathfrak{O}}.$$

Hence, $\tilde{\mathfrak{O}}$ is an $q\mathcal{P}$ -($\in_{\tilde{\sigma}}, \in_{\tilde{\sigma}} \vee q_{\tilde{\tau}}$)- \mathcal{FFI} of $\tilde{\aleph}$.

On the contrary, assume $\tilde{\mathfrak{O}}$ is a $q\mathcal{P}$ -($\in_{\tilde{\sigma}}, \in_{\tilde{\sigma}} \vee q_{\tilde{\tau}}$)- \mathcal{FFI} of $\tilde{\aleph}$. Then \tilde{C} is equal to $\tilde{\mathfrak{O}}_{\tilde{\sigma}}$. Consequently, according to Theorem 4.1, \tilde{C} is a \mathcal{FI} of $\tilde{\aleph}$.

Corollary 1. Let $\tilde{\sigma} + \hat{1} = 2\tilde{\tau}$ and $\phi \neq \tilde{C} \subseteq \tilde{\aleph}$. Then, \tilde{C} is a \mathcal{FI} of $\tilde{\aleph}$ if and only if the characteristic function $\hat{\chi}_{\tilde{C}}$ is a $q\mathcal{P}$ -($\in_{\tilde{\sigma}}, \in_{\tilde{\sigma}} \vee q_{\tilde{\tau}}$)- \mathcal{FFI} of $\tilde{\aleph}$.

Theorem 5. Let $\phi \neq \tilde{C} \subseteq \tilde{\aleph}$ and $\tilde{\sigma} + \hat{1} = 2\tilde{\tau}$. Then, \tilde{C} is a \mathcal{FI} of $\tilde{\aleph}$ if and only if the $q\mathcal{P}$ - \mathcal{F} subset $\tilde{\mathfrak{O}}$ of $\tilde{\aleph}$ defined by the following conditions:

$$(1) \tilde{\mathfrak{O}}(\zeta) \geq \tilde{\tau}, \forall \zeta \in \tilde{C},$$

$$(2) \tilde{\mathfrak{O}}(\zeta) \leq \tilde{\sigma}, \forall \zeta \notin \tilde{C}$$

is a $q\mathcal{P}$ -($q_{\tilde{\tau}}, \in_{\tilde{\sigma}} \vee q_{\tilde{\tau}}$)- \mathcal{FFI} of $\tilde{\aleph}$.

Proof. Let \tilde{C} be a \mathcal{FI} of $\tilde{\aleph}$, $\zeta, \dot{\rho}, \dot{\kappa} \in \tilde{\aleph}$ and let $\tilde{\sigma} < \tilde{\rho}, \tilde{\eta} \leq \hat{1}$ be such that $((\zeta \mathbin{\dot{\vee}} \dot{\rho}) \mathbin{\dot{\vee}} \dot{\kappa})_{\tilde{\rho} q_{\tilde{\tau}} \tilde{\mathfrak{O}}}$ and $\dot{\kappa}_{\tilde{\eta} q_{\tilde{\tau}} \tilde{\mathfrak{O}}}$.

Then

$$\tilde{\mathfrak{O}}(((\zeta \mathbin{\dot{\vee}} \dot{\rho}) \mathbin{\dot{\vee}} \dot{\kappa})) + \tilde{\rho} > 2\tilde{\tau} \Rightarrow \tilde{\mathfrak{O}}((\zeta \mathbin{\dot{\vee}} \dot{\rho}) \mathbin{\dot{\vee}} \dot{\kappa}) > 2\tilde{\tau} - \tilde{\rho} \geq 2\tilde{\tau} - \hat{1} = \tilde{\sigma}$$

and

$$\tilde{\mathfrak{O}}(\dot{\kappa}) + \tilde{\eta} > 2\tilde{\tau} \Rightarrow \tilde{\mathfrak{O}}(\dot{\kappa}) > 2\tilde{\tau} - \tilde{\eta} \geq 2\tilde{\tau} - \hat{1} = \tilde{\sigma}.$$

Thus,

$$\zeta \mathbin{\dot{\vee}} (\dot{\rho} \mathbin{\dot{\vee}} (\dot{\rho} \mathbin{\dot{\vee}} \zeta)) \in \tilde{C} \Rightarrow \tilde{\mathfrak{O}}(\zeta \mathbin{\dot{\vee}} (\dot{\rho} \mathbin{\dot{\vee}} (\dot{\rho} \mathbin{\dot{\vee}} \zeta))) \geq \tilde{\tau}.$$

Now, if $\tilde{\rho} \wedge \tilde{\eta} \leq \tilde{\tau}$, then

$$\tilde{\mathfrak{O}}(\zeta \mathbin{\dot{\vee}} (\dot{\rho} \mathbin{\dot{\vee}} (\dot{\rho} \mathbin{\dot{\vee}} \zeta))) \geq \tilde{\tau} \geq \tilde{\rho} \wedge \tilde{\eta} > \tilde{\sigma}.$$

Hence, $(\zeta \mathbin{\dot{\vee}} (\dot{\rho} \mathbin{\dot{\vee}} (\dot{\rho} \mathbin{\dot{\vee}} \zeta)))_{\tilde{\rho} \wedge \tilde{\eta}} \in_{\tilde{\sigma}} \tilde{\mathfrak{O}}$.

If $\tilde{\rho} \wedge \tilde{\eta} > \tilde{\tau}$, then

$$\tilde{\mathfrak{O}}(\zeta \mathbin{\dot{\vee}} (\dot{\rho} \mathbin{\dot{\vee}} (\dot{\rho} \mathbin{\dot{\vee}} \zeta))) + \tilde{\rho} \wedge \tilde{\eta} > \tilde{\tau} + \tilde{\tau} = 2\tilde{\tau} \Rightarrow (\zeta \mathbin{\dot{\vee}} (\dot{\rho} \mathbin{\dot{\vee}} (\dot{\rho} \mathbin{\dot{\vee}} \zeta)))_{\tilde{\rho} \wedge \tilde{\eta}} q_{\tilde{\tau}} \tilde{\mathfrak{O}}.$$

Therefore, $(\zeta \mathbin{\dot{\vee}} (\dot{\rho} \mathbin{\dot{\vee}} (\dot{\rho} \mathbin{\dot{\vee}} \zeta)))_{\tilde{\rho} \wedge \tilde{\eta}} \in \vee q_{\tilde{\tau}} \tilde{\mathfrak{O}}$. Thus, $\tilde{\mathfrak{O}}$ is an $q\mathcal{P}$ -($q_{\tilde{\tau}}, \in_{\tilde{\sigma}} \vee q_{\tilde{\tau}}$)- \mathcal{FFI} of $\tilde{\aleph}$.

On the contrary, assume $\tilde{\mathfrak{O}}$ is a $q\mathcal{P}$ -($q_{\tilde{\tau}}, \in_{\tilde{\sigma}} \vee q_{\tilde{\tau}}$)- \mathcal{FFI} of $\tilde{\aleph}$. Then, \tilde{C} is equal to $\tilde{\mathfrak{O}}_{\tilde{\sigma}}$. Consequently, according to Theorem 4.1, \tilde{C} is a \mathcal{FI} of $\tilde{\aleph}$.

Corollary 2. Let $\phi \neq \tilde{C} \subseteq \tilde{\aleph}$ and $\tilde{\sigma} + \hat{1} = 2\tilde{\tau}$. Then, \tilde{C} is a \mathcal{FI} of $\tilde{\aleph}$ if and only if the characteristic function $\hat{\chi}_{\tilde{C}}$ is a $q\mathcal{P}$ - $(q_{\tilde{\tau}}, \in_{\tilde{\sigma}} \vee q_{\tilde{\tau}})$ - \mathcal{FFI} of $\tilde{\aleph}$.

Theorem 6. Let $\phi \neq \tilde{C} \subseteq \tilde{\aleph}$ and $\tilde{\sigma} + \hat{1} = 2\tilde{\tau}$. Then, \tilde{C} is a \mathcal{FI} of $\tilde{\aleph}$ if and only if $q\mathcal{P}$ - \mathcal{F} subset $\tilde{\mathcal{D}}$ of $\tilde{\aleph}$ defined by the following conditions:

$$(1) \tilde{\mathcal{D}}(\zeta) \geq \tilde{\tau}, \forall \zeta \in \tilde{C},$$

$$(2) \tilde{\mathcal{D}}(\zeta) \leq \tilde{\sigma}, \forall \zeta \notin \tilde{C}$$

is a $q\mathcal{P}$ - $(\in_{\tilde{\sigma}} \vee q_{\tilde{\tau}}, \in_{\tilde{\sigma}} \vee q_{\tilde{\tau}})$ - \mathcal{FFI} of $\tilde{\aleph}$.

Proof. Let \tilde{C} be a \mathcal{FI} of $\tilde{\aleph}$, $\zeta, \dot{\rho}, \dot{\kappa} \in \tilde{\aleph}$ and let $\tilde{\sigma} < \tilde{\rho}, \tilde{\eta} \leq \hat{1}$ be such that

$$((\zeta \dot{\vee} \dot{\rho}) \dot{\vee} \dot{\kappa})_{\tilde{\rho}} \in_{\tilde{\sigma}} \vee q_{\tilde{\tau}} \tilde{\mathcal{D}} \Rightarrow ((\zeta \dot{\vee} \dot{\rho}) \dot{\vee} \dot{\kappa})_{\tilde{\rho}} \in_{\tilde{\sigma}} \tilde{\mathcal{D}} \text{ or } ((\zeta \dot{\vee} \dot{\rho}) \dot{\vee} \dot{\kappa})_{\tilde{\rho}} q_{\tilde{\tau}} \tilde{\mathcal{D}}$$

and

$$\dot{\kappa}_{\tilde{\eta}} \in_{\tilde{\sigma}} \vee q_{\tilde{\tau}} \tilde{\mathcal{D}} \Rightarrow \dot{\kappa}_{\tilde{\eta}} \in_{\tilde{\sigma}} \tilde{\mathcal{D}} \text{ or } \dot{\kappa}_{\tilde{\eta}} q_{\tilde{\tau}} \tilde{\mathcal{D}}.$$

If $((\zeta \dot{\vee} \dot{\rho}) \dot{\vee} \dot{\kappa})_{\tilde{\rho}} \in_{\tilde{\sigma}} \tilde{\mathcal{D}}$, and $\dot{\kappa}_{\tilde{\eta}} q_{\tilde{\tau}} \tilde{\mathcal{D}}$, then

$$\tilde{\mathcal{D}}((\zeta \dot{\vee} \dot{\rho}) \dot{\vee} \dot{\kappa}) \geq \tilde{\rho} > \tilde{\sigma} \text{ and } \tilde{\mathcal{D}}(\dot{\kappa}) + \tilde{\eta} > 2\tilde{\tau} \Rightarrow \tilde{\mathcal{D}}(\dot{\kappa}) > 2\tilde{\tau} - \tilde{\eta} \geq 2\tilde{\tau} - \hat{1} = \tilde{\sigma}.$$

Thus, $\zeta \dot{\vee} (\dot{\rho} \dot{\vee} (\dot{\rho} \dot{\vee} \dot{\zeta})) \in \tilde{C}$. Analogous as in Theorems 4.3 and 4.5,

$$(\zeta \dot{\vee} (\dot{\rho} \dot{\vee} (\dot{\rho} \dot{\vee} \dot{\zeta})))_{\tilde{\rho} \wedge \tilde{\eta}} \in_{\tilde{\sigma}} \vee q_{\tilde{\tau}} \tilde{\mathcal{D}}.$$

Hence, $\tilde{\mathcal{D}}$ is a $q\mathcal{P}$ - $(\in_{\tilde{\sigma}} \vee q_{\tilde{\tau}}, \in_{\tilde{\sigma}} \vee q_{\tilde{\tau}})$ - \mathcal{FFI} of $\tilde{\aleph}$. The other scenarios can be approached in a similar manner to this one.

On the contrary, assume $\tilde{\mathcal{D}}$ is a $q\mathcal{P}$ - $(\in_{\tilde{\sigma}} \vee q_{\tilde{\tau}}, \in_{\tilde{\sigma}} \vee q_{\tilde{\tau}})$ - \mathcal{FFI} of $\tilde{\aleph}$. Then, \tilde{C} is equal to $\tilde{\mathcal{D}}_{\tilde{\sigma}}$. Consequently, according to Theorem 4.1, \tilde{C} is a \mathcal{FI} of $\tilde{\aleph}$.

Corollary 3. Let $\phi \neq \tilde{C} \subseteq \tilde{\aleph}$ and $\tilde{\sigma} + \hat{1} = 2\tilde{\tau}$. Then, \tilde{C} is a \mathcal{FI} of $\tilde{\aleph}$ if and only if the characteristic function $\hat{\chi}_{\tilde{C}}$ is a $q\mathcal{P}$ - $(\in_{\tilde{\sigma}} \vee q_{\tilde{\tau}}, \in_{\tilde{\sigma}} \vee q_{\tilde{\tau}})$ - \mathcal{FFI} of $\tilde{\aleph}$.

Theorem 7. Every $q\mathcal{P}$ - $(q_{\tilde{\tau}}, \in_{\tilde{\sigma}} \vee q_{\tilde{\tau}})$ - \mathcal{FFI} of $\tilde{\aleph}$ is a $q\mathcal{P}$ - $(\in_{\tilde{\sigma}}, \in_{\tilde{\sigma}} \vee q_{\tilde{\tau}})$ - \mathcal{FFI} of $\tilde{\aleph}$.

Proof. Let $\tilde{\mathcal{D}}$ be a $q\mathcal{P}$ - $(q_{\tilde{\tau}}, \in_{\tilde{\sigma}} \vee q_{\tilde{\tau}})$ - \mathcal{FFI} of $\tilde{\aleph}$, $\zeta, \dot{\rho}, \dot{\kappa} \in \tilde{\aleph}$ and let $\tilde{\sigma} < \tilde{\rho}, \tilde{\eta} \leq \hat{1}$ be such that $\zeta \dot{\vee} (\dot{\rho} \dot{\vee} (\dot{\rho} \dot{\vee} \dot{\zeta}))_{\tilde{\rho}} \in_{\tilde{\sigma}} \tilde{\mathcal{D}}$ and $\dot{\kappa}_{\tilde{\eta}} \in_{\tilde{\tau}} \tilde{\mathcal{D}}$. Then,

$$\tilde{\mathcal{D}}(\zeta \dot{\vee} (\dot{\rho} \dot{\vee} (\dot{\rho} \dot{\vee} \dot{\zeta}))) \geq \tilde{\rho} > \tilde{\sigma} \text{ and } \tilde{\mathcal{D}}(\dot{\kappa}) \geq \tilde{\eta} > \tilde{\sigma}.$$

Suppose that $(\zeta \dot{\vee} (\dot{\rho} \dot{\vee} (\dot{\rho} \dot{\vee} \dot{\zeta})))_{\tilde{\rho} \wedge \tilde{\eta}} \in_{\tilde{\sigma}} \overline{\vee q_{\tilde{\tau}} \tilde{\mathcal{D}}}$. Then

$$\tilde{\mathcal{D}}(\zeta \dot{\vee} (\dot{\rho} \dot{\vee} (\dot{\rho} \dot{\vee} \dot{\zeta}))) + \tilde{\mathcal{D}}(\zeta \dot{\vee} (\dot{\rho} \dot{\vee} (\dot{\rho} \dot{\vee} \dot{\zeta}))) < \tilde{\mathcal{D}}(\zeta \dot{\vee} (\dot{\rho} \dot{\vee} (\dot{\rho} \dot{\vee} \dot{\zeta}))) + \tilde{\rho} \wedge \tilde{\varpi} \leq 2\tilde{\tau}.$$

Therefore, $\tilde{\mathcal{D}}(\zeta \dot{\vee} (\dot{\rho} \dot{\vee} (\dot{\rho} \dot{\vee} \dot{\zeta}))) < \tilde{\tau}$.

Now,

$$\tilde{\mathcal{D}}(\zeta \dot{\vee} (\dot{\rho} \dot{\vee} (\dot{\rho} \dot{\vee} \dot{\zeta}))) \vee \tilde{\sigma} < \tilde{\mathcal{D}}((\zeta \dot{\vee} \dot{\rho}) \dot{\vee} \dot{\kappa}) \wedge \tilde{\tau}.$$

Choose $\tilde{\sigma} < \hat{r} \leq \hat{1}$ Then

$$2\tilde{\tau} - \tilde{\mathfrak{D}}(\zeta \tilde{\mathfrak{J}} (\dot{\rho} \tilde{\mathfrak{J}} (\dot{\rho} \tilde{\mathfrak{J}} \zeta))) \vee \tilde{\sigma} \geq \hat{r} > 2\tilde{\tau} - \tilde{\mathfrak{D}}((\zeta \tilde{\mathfrak{J}} \dot{\rho}) \wedge \tilde{\mathfrak{D}}(\dot{\kappa}) \wedge \tilde{\tau}.$$

Therefore,

$$2\tilde{\tau} - \tilde{\mathfrak{D}}(\zeta \tilde{\mathfrak{J}} (\dot{\rho} \tilde{\mathfrak{J}} (\dot{\rho} \tilde{\mathfrak{J}} \zeta))) \wedge (2\tilde{\tau} - \tilde{\sigma}) \geq (2\tilde{\tau} - \tilde{\mathfrak{D}}((\zeta \tilde{\mathfrak{J}} \dot{\rho})) \vee (2\tilde{\tau} - \tilde{\mathfrak{D}}(\dot{\kappa})) \vee \tilde{\tau}.$$

Thus,

$$\hat{r} > 2\tilde{\tau} - \tilde{\mathfrak{D}}((\zeta \tilde{\mathfrak{J}} \dot{\rho}), \hat{r} > 2\tilde{\tau} - \tilde{\mathfrak{D}}(\dot{\kappa}) \text{ and } 2\tilde{\tau} - \tilde{\mathfrak{D}}(\zeta \tilde{\mathfrak{J}} (\dot{\rho} \tilde{\mathfrak{J}} (\dot{\rho} \tilde{\mathfrak{J}} \zeta))) > \hat{r}$$

So,

$$\tilde{\mathfrak{D}}((\zeta \tilde{\mathfrak{J}} \dot{\rho}) + \hat{r} > 2\tilde{\tau}, \tilde{\mathfrak{D}}(\dot{\kappa}) + \hat{r} > 2\tilde{\tau} \text{ and } \tilde{\mathfrak{D}}(\zeta \tilde{\mathfrak{J}} (\dot{\rho} \tilde{\mathfrak{J}} (\dot{\rho} \tilde{\mathfrak{J}} \zeta))) + \hat{r} < 2\tilde{\tau}.$$

Thus, $\zeta \tilde{\mathfrak{J}} (\dot{\rho} \tilde{\mathfrak{J}} (\dot{\rho} \tilde{\mathfrak{J}} \zeta))_{\hat{r}} q_{\tilde{\tau}} \tilde{\mathfrak{D}}, \dot{\kappa}_{\hat{r}} q_{\tilde{\tau}} \tilde{\mathfrak{D}}$ but $(\zeta \tilde{\mathfrak{J}} (\dot{\rho} \tilde{\mathfrak{J}} (\dot{\rho} \tilde{\mathfrak{J}} \zeta)))_{\hat{r}} \in_{\tilde{\sigma}} \vee q_{\tilde{\tau}} \tilde{\mathfrak{D}}$, a contradiction. Hence, $\tilde{\mathfrak{D}}$ is a $q\mathcal{P}$ -($\in_{\tilde{\sigma}}, \in_{\tilde{\sigma}} \vee q_{\tilde{\tau}}$)- \mathcal{FFI} of $\tilde{\aleph}$.

Theorem 8. Every $q\mathcal{P}$ -($\in_{\tilde{\sigma}} \vee q_{\tilde{\tau}}, \in_{\tilde{\sigma}} \vee q_{\tilde{\tau}}$)- \mathcal{FFI} of $\tilde{\aleph}$ is a $q\mathcal{P}$ -($\in_{\tilde{\sigma}}, \in_{\tilde{\sigma}} \vee q_{\tilde{\tau}}$)- \mathcal{FFI} of $\tilde{\aleph}$.

Proof. The proof is based on the observation that if $\zeta_{\tilde{\rho}} \in_{\tilde{\sigma}} \tilde{\mathfrak{D}}$, then $\zeta_{\tilde{\rho}} \in_{\tilde{\sigma}} \vee q_{\tilde{\tau}} \tilde{\mathfrak{D}}$.

Theorem 9. Every $q\mathcal{P}$ -($\in_{\tilde{\sigma}}, \in_{\tilde{\sigma}}$)- \mathcal{FFI} is a $q\mathcal{P}$ -($\in_{\tilde{\sigma}}, \in_{\tilde{\sigma}} \vee q_{\tilde{\tau}}$)- \mathcal{FFI} of $\tilde{\aleph}$.

Proof. Let $\tilde{\mathfrak{D}}$ be a $q\mathcal{P}$ -($\in_{\tilde{\sigma}}, \in_{\tilde{\sigma}}$)- \mathcal{FFI} of $\tilde{\aleph}$, $\zeta, \dot{\rho}, \dot{\kappa} \in \tilde{\aleph}$ and let $\tilde{\sigma} < \tilde{\rho}, \tilde{\eta} \leq \hat{1}$. So the $q\mathcal{P}$ - \mathcal{F} points $((\zeta \tilde{\mathfrak{J}} \dot{\rho}) \tilde{\mathfrak{J}} \dot{\kappa})_{\tilde{\rho}} \in_{\tilde{\sigma}} \tilde{\mathfrak{D}}$ and $\dot{\kappa}_{\tilde{\eta}} \in_{\tilde{\sigma}} \tilde{\mathfrak{D}}$. Then

$$((\zeta \tilde{\mathfrak{J}} \dot{\rho}) \tilde{\mathfrak{J}} \dot{\kappa})_{\tilde{\rho}} \in_{\tilde{\sigma}} \tilde{\mathfrak{D}} \text{ and } \dot{\kappa}_{\tilde{\eta}} \in_{\tilde{\sigma}} \tilde{\mathfrak{D}} \Rightarrow (\zeta \tilde{\mathfrak{J}} (\dot{\rho} \tilde{\mathfrak{J}} (\dot{\rho} \tilde{\mathfrak{J}} \zeta)))_{\tilde{\rho} \wedge \tilde{\eta}} \in_{\tilde{\sigma}} \vee q_{\tilde{\tau}} \tilde{\mathfrak{D}}.$$

Thus, $\tilde{\mathfrak{D}}$ is a $q\mathcal{P}$ -($\in_{\tilde{\sigma}}, \in_{\tilde{\sigma}} \vee q_{\tilde{\tau}}$)- \mathcal{FFI} of $\tilde{\aleph}$.

5. Quadri-Polar ($\in_{\tilde{\sigma}}, \in_{\tilde{\sigma}} \vee q_{\tilde{\tau}}$)-Fuzzy Fantastic Ideals

In this section, we present the notion of a $q\mathcal{P}$ -($\in_{\tilde{\sigma}}, \in_{\tilde{\sigma}} \vee q_{\tilde{\tau}}$)- $\mathcal{FFI}(s)$ and explore several of its essential characteristics and properties.

Definition 5. A $q\mathcal{P}$ - \mathcal{F} set $\tilde{\mathfrak{D}}$ of $\tilde{\aleph}$ is a $q\mathcal{P}$ -($\in_{\tilde{\sigma}}, \in_{\tilde{\sigma}} \vee q_{\tilde{\tau}}$)- \mathcal{FFI} of $\tilde{\aleph}$ if it satisfies condition (1), as follows:

(1) $((\zeta \tilde{\mathfrak{J}} \dot{\rho}) \tilde{\mathfrak{J}} \dot{\kappa})_{\tilde{\rho}} \in_{\tilde{\sigma}} \tilde{\mathfrak{D}}, \dot{\kappa}_{\tilde{\eta}} \in_{\tilde{\sigma}} \tilde{\mathfrak{D}} \Rightarrow (\zeta \tilde{\mathfrak{J}} (\dot{\rho} \tilde{\mathfrak{J}} (\dot{\rho} \tilde{\mathfrak{J}} \zeta)))_{\tilde{\rho} \wedge \tilde{\eta}} \in_{\tilde{\sigma}} \vee q_{\tilde{\tau}} \tilde{\mathfrak{D}}, \forall \tilde{\sigma} < \tilde{\rho}, \tilde{\eta} \leq \hat{1}$ and $\zeta, \dot{\rho}, \dot{\kappa} \in \tilde{\aleph}$.

Example 2. Consider the BCK-algebra $(\tilde{\aleph}; \tilde{\mathfrak{J}}, 0)$ and a $q\mathcal{P}$ - \mathcal{F} set $\tilde{\mathfrak{D}}$ as illustrated in Example 3.1. It is evident from Definition 5.1, $\tilde{\mathfrak{D}}$ is a $q\mathcal{P}$ -($\in_{(0.2, 0.1, 0.3, 0.2)}, \in_{(0.2, 0.1, 0.3, 0.2)} \vee q_{(0.61, 0.68, 0.78, 0.57)}$)- \mathcal{FFI} of $\tilde{\aleph}$.

Theorem 10. For a $q\mathcal{P}$ - \mathcal{F} set $\tilde{\mathfrak{D}}$ of $\tilde{\aleph}$, condition (1) in Definition 5.1 is similar with condition (2), as follows:

(2) $\tilde{\mathfrak{D}}(\zeta \tilde{\mathfrak{J}} (\dot{\rho} \tilde{\mathfrak{J}} (\dot{\rho} \tilde{\mathfrak{J}} \zeta))) \vee \tilde{\sigma} \geq \tilde{\mathfrak{D}}((\zeta \tilde{\mathfrak{J}} \dot{\rho}) \tilde{\mathfrak{J}} \dot{\kappa}) \wedge \tilde{\mathfrak{D}}(\dot{\kappa}), \tilde{\tau}, \forall \zeta, \dot{\rho} \in \tilde{\aleph}$.

Proof. (1) \Rightarrow (2). Assume that (2) does not hold. Then, $\exists \zeta, \varrho, \kappa \in \tilde{\mathfrak{N}}$ such that

$$\tilde{\partial}(\zeta \tilde{\vee} (\varrho \tilde{\vee} (\varrho \tilde{\vee} \zeta))) \vee \tilde{\sigma} < \tilde{\partial}((\zeta \tilde{\vee} \varrho) \tilde{\vee} \kappa) \wedge \tilde{\partial}(\kappa) \wedge \tilde{\tau}.$$

Then,

$$\tilde{\partial}(\zeta \tilde{\vee} (\varrho \tilde{\vee} (\varrho \tilde{\vee} \zeta))) \vee \tilde{\sigma} < \tilde{\rho} \leq \tilde{\partial}((\zeta \tilde{\vee} \varrho) \tilde{\vee} \kappa) \wedge \tilde{\partial}(\kappa) \wedge \tilde{\tau}.$$

Thus,

$$((\zeta \tilde{\vee} \varrho) \tilde{\vee} \kappa)_{\tilde{\rho}} \in_{\tilde{\sigma}} \tilde{\partial} \text{ and } \kappa_{\tilde{\rho}} \in_{\tilde{\sigma}} \tilde{\partial}.$$

But $(\zeta \tilde{\vee} (\varrho \tilde{\vee} (\varrho \tilde{\vee} \zeta)))_{\tilde{\rho} \in_{\tilde{\sigma}} \tilde{\partial} \vee \in_{\tilde{\sigma}} \tilde{\partial}}$, a contradiction.

$$\tilde{\partial}(\zeta \tilde{\vee} (\varrho \tilde{\vee} (\varrho \tilde{\vee} \zeta))) \vee \tilde{\sigma} \geq \tilde{\partial}((\zeta \tilde{\vee} \varrho) \tilde{\vee} \kappa) \wedge \tilde{\partial}(\kappa) \wedge \tilde{\tau}.$$

(2) \Rightarrow (1) Let $((\zeta \tilde{\vee} \varrho) \tilde{\vee} \kappa)_{\tilde{\rho}} \in_{\tilde{\sigma}} \tilde{\partial}, \kappa_{\tilde{\eta}} \in_{\tilde{\sigma}} \tilde{\partial}$. Then, $\tilde{\partial}((\zeta \tilde{\vee} \varrho) \tilde{\vee} \kappa) \geq \tilde{\rho} > \tilde{\sigma}$ and $\tilde{\partial}(\kappa) \geq \tilde{\eta} > \tilde{\sigma}$. If $(\zeta \tilde{\vee} (\varrho \tilde{\vee} (\varrho \tilde{\vee} \zeta)))_{\tilde{\rho} \wedge \tilde{\eta}} \in_{\tilde{\sigma}}$, then (1) is hold.

If $(\zeta \tilde{\vee} (\varrho \tilde{\vee} (\varrho \tilde{\vee} \zeta)))_{\tilde{\rho} \wedge \tilde{\eta}} \in_{\tilde{\sigma}} \tilde{\partial}$, then $\tilde{\partial}(\zeta \tilde{\vee} (\varrho \tilde{\vee} (\varrho \tilde{\vee} \zeta))) < \tilde{\rho} \wedge \tilde{\eta}$. Since

$$\begin{aligned} \tilde{\partial}(\zeta \tilde{\vee} (\varrho \tilde{\vee} (\varrho \tilde{\vee} \zeta))) \vee \tilde{\sigma} &\geq \tilde{\partial}(\zeta) \wedge \tilde{\partial}(\varrho) \wedge \tilde{\tau} \\ &\geq \tilde{\rho} \wedge \tilde{\omega} \wedge \tilde{\tau}. \end{aligned}$$

Therefore,

$$\tilde{\partial}(\zeta \tilde{\vee} (\varrho \tilde{\vee} (\varrho \tilde{\vee} \zeta))) \geq \tilde{\tau} \text{ and } \tilde{\rho} \wedge \tilde{\omega} > \tilde{\tau}.$$

Thus,

$$\tilde{\partial}(\zeta \tilde{\vee} (\varrho \tilde{\vee} (\varrho \tilde{\vee} \zeta))) + \tilde{\rho} \wedge \tilde{\eta} > \tilde{\tau} + \tilde{\tau} = 2\tilde{\tau} \Rightarrow (\zeta \tilde{\vee} (\varrho \tilde{\vee} (\varrho \tilde{\vee} \zeta)))_{\tilde{\rho} \wedge \tilde{\eta} q_{\tilde{\tau}}} \tilde{\partial}.$$

Hence, $(\zeta \tilde{\vee} (\varrho \tilde{\vee} (\varrho \tilde{\vee} \zeta)))_{\tilde{\rho} \wedge \tilde{\eta}} \in_{\tilde{\sigma}} \vee q_{\tilde{\tau}} \tilde{\partial}$.

Corollary 4. A $q\mathcal{P}$ - \mathcal{F} set $\tilde{\partial}$ of $\tilde{\mathfrak{N}}$ is a $q\mathcal{P}$ -($\in_{\tilde{\sigma}}, \in_{\tilde{\sigma}} \vee q_{\tilde{\tau}}$)- \mathcal{FFI} of $\tilde{\mathfrak{N}}$ if it satisfies $\tilde{\partial}(\zeta \tilde{\vee} (\varrho \tilde{\vee} (\varrho \tilde{\vee} \zeta))) \vee \tilde{\sigma} \geq \tilde{\partial}((\zeta \tilde{\vee} \varrho) \tilde{\vee} \kappa) \wedge \tilde{\partial}(\kappa), \tilde{\tau}, \forall \zeta, \varrho \in \tilde{\mathfrak{N}}$.

Theorem 11. The intersection of any collection of $q\mathcal{P}$ -($\in_{\tilde{\sigma}}, \in_{\tilde{\sigma}} \vee q_{\tilde{\tau}}$)- \mathcal{FFI} s of $\tilde{\mathfrak{N}}$ is a $q\mathcal{P}$ -($\in_{\tilde{\sigma}}, \in_{\tilde{\sigma}} \vee q_{\tilde{\tau}}$)- \mathcal{FFI} of $\tilde{\mathfrak{N}}$.

Proof. Let $\{\tilde{\partial}_i\}_{i \in I}$ be a collection of $q\mathcal{P}$ -($\in_{\tilde{\sigma}}, \in_{\tilde{\sigma}} \vee q_{\tilde{\tau}}$)- \mathcal{FFI} s of $\tilde{\mathfrak{N}}$ and $(\zeta \tilde{\vee} \varrho) \tilde{\vee} \kappa, \kappa \in \tilde{\mathfrak{N}}$. Then,

$$\tilde{\partial}_i(\zeta \tilde{\vee} (\varrho \tilde{\vee} (\varrho \tilde{\vee} \zeta))) \vee \tilde{\sigma} \geq \tilde{\partial}_i((\zeta \tilde{\vee} \varrho) \tilde{\vee} \kappa) \wedge \tilde{\partial}_i(\kappa) \wedge \tilde{\tau}.$$

Thus,

$$\begin{aligned} (\wedge_{i \in I} \tilde{\partial}_i)(\zeta \tilde{\vee} (\varrho \tilde{\vee} (\varrho \tilde{\vee} \zeta))) \vee \tilde{\sigma} &= \wedge_{i \in I} \tilde{\partial}_i(\zeta \tilde{\vee} (\varrho \tilde{\vee} (\varrho \tilde{\vee} \zeta))) \vee \tilde{\sigma} \\ &\geq \wedge_{i \in I} (\tilde{\partial}_i((\zeta \tilde{\vee} \varrho) \tilde{\vee} \kappa) \wedge \tilde{\partial}_i(\kappa) \wedge \tilde{\tau}) \\ &\geq (\wedge_{i \in I} \tilde{\partial}_i)((\zeta \tilde{\vee} \varrho) \tilde{\vee} \kappa) \wedge (\wedge_{i \in I} \tilde{\partial}_i)(\kappa) \wedge \tilde{\tau}. \end{aligned}$$

Therefore, $(\wedge_{i \in I} \tilde{\mathfrak{d}}_i)(\zeta \check{\vee} (\varrho \check{\vee} (\varrho \check{\vee} \zeta))) \vee \tilde{\sigma} \geq (\wedge_{i \in I} \tilde{\mathfrak{d}}_i)((\zeta \check{\vee} \varrho) \check{\vee} \kappa) \wedge (\wedge_{i \in I} \tilde{\mathfrak{d}}_i)(\kappa) \wedge \tilde{\tau}$. Hence, $\wedge_{i \in I} \tilde{\mathfrak{d}}_i$ is a $q\mathcal{P}-(\in_{\tilde{\sigma}}, \in_{\tilde{\sigma}} \vee q_{\tilde{\tau}})\mathcal{FFI}$ of $\tilde{\aleph}$.

For any $q\mathcal{P}\text{-}\mathcal{F}$ set $\tilde{\mathfrak{d}}$ of $\tilde{\aleph}$ and $\tilde{\rho} \in [0, 1]^q$, we define:

- (1) $\tilde{\mathfrak{d}}_{\tilde{\rho}}^{\tilde{\sigma}} = \{\zeta \in \tilde{\aleph} \mid \zeta_{\tilde{\rho}} \in_{\tilde{\sigma}} \tilde{\mathfrak{d}}\}$,
- (2) $\langle \tilde{\mathfrak{d}} \rangle_{\tilde{\rho}}^{\tilde{\tau}} = \{\zeta \in \tilde{\aleph} \mid \zeta_{\tilde{\rho}} q_{\tilde{\tau}} \tilde{\mathfrak{d}}\}$,
- (3) $[\tilde{\mathfrak{d}}]_{\tilde{\rho}}^{\tilde{\tau}} = \{\zeta \in \tilde{\aleph} \mid \zeta_{\tilde{\rho}} \in_{\tilde{\sigma}} \vee q_{\tilde{\tau}} \tilde{\mathfrak{d}}\}$.

It is clear that

$$[\tilde{\mathfrak{d}}]_{\tilde{\rho}}^{\tilde{\tau}} = \tilde{\mathfrak{d}}_{\tilde{\rho}}^{\tilde{\sigma}} \cup \langle \tilde{\mathfrak{d}} \rangle_{\tilde{\rho}}^{\tilde{\tau}}.$$

The ensuing theorems elucidate the connection between $q\mathcal{P}-(\in_{\tilde{\sigma}}, \in_{\tilde{\sigma}} \vee q_{\tilde{\tau}})\mathcal{FFIs}$ and the crisp \mathcal{FI} s in $\tilde{\aleph}$.

Theorem 12. *Let $\tilde{\mathfrak{d}}$ be a $q\mathcal{P}\text{-}\mathcal{F}$ set of $\tilde{\aleph}$. Then, $\tilde{\mathfrak{d}}$ is a $q\mathcal{P}-(\in_{\tilde{\sigma}}, \in_{\tilde{\sigma}} \vee q_{\tilde{\tau}})\mathcal{FFI}$ of $\tilde{\aleph} \Leftrightarrow \tilde{\mathfrak{d}}_{\tilde{\rho}}^{\tilde{\sigma}} \neq \emptyset$ is a \mathcal{FI} of $\tilde{\aleph}$, $\forall \tilde{\sigma} < \tilde{\rho} \leq \tilde{\tau}$.*

Proof. Let $\tilde{\mathfrak{d}}$ be a $q\mathcal{P}-(\in_{\tilde{\sigma}}, \in_{\tilde{\sigma}} \vee q_{\tilde{\tau}})\mathcal{FFI}$ of $\tilde{\aleph}$ and let $\zeta, \varrho, \kappa \in \tilde{\mathfrak{d}}_{\tilde{\rho}}^{\tilde{\sigma}}$ for $\tilde{\sigma} < \tilde{\rho} \leq \tilde{\tau}$. Then $\tilde{\mathfrak{d}}((\zeta \check{\vee} \varrho) \check{\vee} \kappa) \geq \tilde{\rho} > \tilde{\sigma}$ and $\tilde{\mathfrak{d}}(\kappa) \geq \tilde{\rho} > \tilde{\sigma}$. Thus, we have

$$\begin{aligned} \tilde{\mathfrak{d}}(\zeta \check{\vee} (\varrho \check{\vee} (\varrho \check{\vee} \zeta))) \vee \tilde{\sigma} &\geq \tilde{\mathfrak{d}}((\zeta \check{\vee} \varrho) \check{\vee} \kappa) \wedge \tilde{\mathfrak{d}}(\kappa) \wedge \tilde{\tau} \\ &\geq \tilde{\rho} \wedge \tilde{\rho} \wedge \tilde{\tau} \\ &= \tilde{\rho} \wedge \tilde{\tau} \\ &= \tilde{\rho} > \tilde{\sigma}. \end{aligned}$$

Therefore,

$$\tilde{\mathfrak{d}}(\zeta \check{\vee} (\varrho \check{\vee} (\varrho \check{\vee} \zeta))) \geq \tilde{\rho} \Rightarrow \zeta \check{\vee} (\varrho \check{\vee} (\varrho \check{\vee} \zeta)) \in \tilde{\mathfrak{d}}_{\tilde{\rho}}^{\tilde{\sigma}}.$$

Thus, $\tilde{\mathfrak{d}}_{\tilde{\rho}}^{\tilde{\sigma}}$ is a \mathcal{FI} of U .

On the other hand, suppose that $\tilde{\mathfrak{d}}_{\tilde{\rho}}^{\tilde{\sigma}}$ is a \mathcal{FI} of U , $\forall \tilde{\sigma} < \tilde{\rho} \leq \tilde{\tau}$. Assume $\zeta, \varrho, \kappa \in \tilde{\aleph}$ such that

$$\tilde{\mathfrak{d}}(\zeta \check{\vee} (\varrho \check{\vee} (\varrho \check{\vee} \zeta))) \vee \tilde{\sigma} < \tilde{\mathfrak{d}}((\zeta \check{\vee} \varrho) \check{\vee} \kappa) \wedge \tilde{\mathfrak{d}}(\kappa) \wedge \tilde{\tau}.$$

Select $\tilde{\sigma} < \tilde{\rho} \leq \tilde{\tau}$ such that

$$\tilde{\mathfrak{d}}(\zeta \check{\vee} (\varrho \check{\vee} (\varrho \check{\vee} \zeta))) \vee \tilde{\sigma} < \hat{r} = \tilde{\mathfrak{d}}((\zeta \check{\vee} \varrho) \check{\vee} \kappa) \wedge \tilde{\mathfrak{d}}(\kappa) \wedge \tilde{\tau}.$$

Then, $((\zeta \check{\vee} \varrho) \check{\vee} \kappa)_{\tilde{\rho}} \in_{\tilde{\sigma}} \tilde{\mathfrak{d}}$, $\kappa_{\tilde{\rho}} \in_{\tilde{\sigma}} \tilde{\mathfrak{d}}$, but $(\zeta \check{\vee} (\varrho \check{\vee} (\varrho \check{\vee} \zeta)))_{\hat{r}} \notin_{\tilde{\sigma}} \vee q_{\tilde{\tau}} \tilde{\mathfrak{d}}$. Since $\tilde{\mathfrak{d}}_{\tilde{\rho}}^{\tilde{\sigma}}$ is a \mathcal{FI} of $\tilde{\aleph}$,

$$\zeta \check{\vee} (\varrho \check{\vee} (\varrho \check{\vee} \zeta)) \in \tilde{\mathfrak{d}}_{\tilde{\rho}}^{\tilde{\sigma}}, \text{ a contradiction.}$$

Hence, $\tilde{\mathfrak{d}}(\zeta \check{\vee} (\varrho \check{\vee} (\varrho \check{\vee} \zeta))) \vee \tilde{\sigma} \geq \tilde{\mathfrak{d}}((\zeta \check{\vee} \varrho) \check{\vee} \kappa) \wedge \tilde{\mathfrak{d}}(\kappa) \wedge \tilde{\tau}$. Therefore, $\tilde{\mathfrak{d}}$ is a $q\mathcal{P}-(\in_{\tilde{\sigma}}, \in_{\tilde{\sigma}} \vee q_{\tilde{\tau}})\mathcal{FFI}$ of $\tilde{\aleph}$.

By setting $\tilde{\sigma} = \tilde{0}$ and $\tilde{\tau} = \widehat{0.5}$ in Theorem 5.3, we can derive the subsequent corollary.

Corollary 5. Let $\tilde{\mathcal{O}}$ be a $q\mathcal{P}$ - \mathcal{F} set of $\tilde{\mathcal{N}}$. Then $\tilde{\mathcal{O}}$ is a $q\mathcal{P}$ -($\in, \in \vee q$) \mathcal{FFI} of $\tilde{\mathcal{N}} \Leftrightarrow \tilde{\mathcal{O}}_{\tilde{\rho}} = \{\dot{\varsigma} \in \tilde{\mathcal{N}} \mid \dot{\varsigma}_{\tilde{\rho}} \in \tilde{\mathcal{O}}\} \neq \phi$ is a \mathcal{FI} of $\tilde{\mathcal{N}}$, $\forall \tilde{\rho} \in (0, 0.5]^q$.

Theorem 13. Let $\tilde{\mathcal{O}}$ be a $q\mathcal{P}$ - \mathcal{F} set of $\tilde{\mathcal{N}}$. Then

- (1) $\tilde{\mathcal{O}}$ is a $q\mathcal{P}$ -($\in_{\tilde{\sigma}}, \in_{\tilde{\sigma}} \vee q_{\tilde{\tau}}$)- \mathcal{FFI} s of $\tilde{\mathcal{N}} \Leftrightarrow \tilde{\mathcal{O}}_{\tilde{\rho}}^{\tilde{\sigma}} \neq \phi$ is a \mathcal{FI} of $\tilde{\mathcal{N}}$, $\forall \tilde{\sigma} < \tilde{\rho} \leq \tilde{\tau}$.
- (2) If $\hat{1} + \tilde{\sigma} = 2\tilde{\tau}$, then $\tilde{\mathcal{O}}$ is a $q\mathcal{P}$ -($\in_{\tilde{\sigma}}, \in_{\tilde{\sigma}} \vee q_{\tilde{\tau}}$)- \mathcal{FFI} of $\tilde{\mathcal{N}} \Leftrightarrow \langle \tilde{\mathcal{O}} \rangle_{\tilde{\rho}}^{\tilde{\tau}} \neq \phi$ is a \mathcal{FI} of $\tilde{\mathcal{N}}$, $\forall \tilde{\tau} < \tilde{\rho} \leq \hat{1}$.
- (3) If $\hat{1} + \tilde{\sigma} = 2\tilde{\tau}$, then $\tilde{\mathcal{O}}$ is a $q\mathcal{P}$ -($\in_{\tilde{\sigma}}, \in_{\tilde{\sigma}} \vee q_{\tilde{\tau}}$)- \mathcal{FFI} of $\tilde{\mathcal{N}} \Leftrightarrow [\tilde{\mathcal{O}}]_{\tilde{\rho}}^{\tilde{\tau}} \neq \phi$ is a \mathcal{FI} of $\tilde{\mathcal{N}}$, $\forall \tilde{\sigma} < \tilde{\rho} \leq \hat{1}$.

Proof. (1) Let $\tilde{\mathcal{O}}$ be a $q\mathcal{P}$ -($\in_{\tilde{\sigma}}, \in_{\tilde{\sigma}} \vee q_{\tilde{\tau}}$)- \mathcal{FFI} of $\tilde{\mathcal{N}}$. Let $\dot{\varsigma} \in \tilde{\mathcal{O}}_{\tilde{\rho}}^{\tilde{\tau}}$. Then

$$\tilde{\mathcal{O}}(0) \vee \tilde{\sigma} \geq \tilde{\mathcal{O}}(\dot{\varsigma}) \wedge \tilde{\tau} \geq \tilde{\rho} \wedge \tilde{\tau} > \tilde{\sigma}.$$

Hence, $\tilde{\mathcal{O}}(0) \geq \tilde{\rho} \Rightarrow 0 \in \tilde{\mathcal{O}}_{\tilde{\rho}}^{\tilde{\tau}}$.

Let $\dot{\varsigma}, \dot{\varrho}, \dot{\kappa} \in \tilde{\mathcal{O}}_{\tilde{\rho}}^{\tilde{\tau}}$. Then,

$$\tilde{\mathcal{O}}((\dot{\varsigma} \tilde{\vee} \dot{\varrho}) \tilde{\vee} \dot{\kappa}) \geq \tilde{\rho} > \tilde{\sigma} \text{ and } \tilde{\mathcal{O}}(\dot{\kappa}) \geq \tilde{\rho} > \tilde{\sigma}.$$

By Theorem 5.1 (2), we have

$$\begin{aligned} \tilde{\mathcal{O}}(\dot{\varsigma} \tilde{\vee} (\dot{\varrho} \tilde{\vee} (\dot{\varrho} \tilde{\vee} \dot{\varsigma}))) \vee \tilde{\sigma} &\geq \tilde{\mathcal{O}}((\dot{\varsigma} \tilde{\vee} \dot{\varrho}) \tilde{\vee} \dot{\kappa}) \wedge \tilde{\tau} \\ &\geq \tilde{\rho} \wedge \tilde{\rho} \wedge \tilde{\tau} \\ &= \tilde{\rho} \wedge \tilde{\tau} \\ &= \tilde{\rho} > \tilde{\sigma}. \end{aligned}$$

Therefore,

$$\tilde{\mathcal{O}}(\dot{\varsigma} \tilde{\vee} (\dot{\varrho} \tilde{\vee} (\dot{\varrho} \tilde{\vee} \dot{\varsigma}))) \geq \tilde{\rho} \Rightarrow \dot{\varsigma} \tilde{\vee} (\dot{\varrho} \tilde{\vee} (\dot{\varrho} \tilde{\vee} \dot{\varsigma})) \in \tilde{\mathcal{O}}_{\tilde{\rho}}^{\tilde{\sigma}}.$$

Hence, $\tilde{\mathcal{O}}_{\tilde{\rho}}^{\tilde{\sigma}}$ is a \mathcal{FI} of $\tilde{\mathcal{N}}$.

Conversely, assume $\tilde{\mathcal{O}}_{\tilde{\rho}}^{\tilde{\sigma}}$ is a \mathcal{FI} of $\tilde{\mathcal{N}}$, $\forall \tilde{\rho} \in (\sigma, \tau]$. Let $\dot{\varsigma} \in \tilde{\mathcal{N}}$ be such that $\tilde{\mathcal{O}}(0) \vee \sigma < \tilde{\rho} = \tilde{\mathcal{O}}(\dot{\varsigma}) \wedge \tilde{\tau}$. Then $\dot{\varsigma}_{\tilde{\rho}} \in_{\tilde{\sigma}} \tilde{\mathcal{O}}$, but $0_{\tilde{\rho}} \in_{\tilde{\sigma}} \overline{\vee q_{\tilde{\tau}}} \tilde{\mathcal{O}}$, a contradiction. Suppose $\dot{\varsigma}, \dot{\varrho}, \dot{\kappa} \in \tilde{\mathcal{N}}$. Then

$$\tilde{\mathcal{O}}(\dot{\varsigma} \tilde{\vee} (\dot{\varrho} \tilde{\vee} (\dot{\varrho} \tilde{\vee} \dot{\varsigma}))) \vee \tilde{\sigma} < \tilde{\mathcal{O}}((\dot{\varsigma} \tilde{\vee} \dot{\varrho}) \tilde{\vee} \dot{\kappa}) \wedge \tilde{\mathcal{O}}(\dot{\kappa}) \wedge \tilde{\tau}.$$

Select some $\tilde{\rho} \in (\tilde{\sigma}, \tilde{\tau}]$ such that

$$\tilde{\mathcal{O}}(\dot{\varsigma} \tilde{\vee} (\dot{\varrho} \tilde{\vee} (\dot{\varrho} \tilde{\vee} \dot{\varsigma}))) \vee \tilde{\sigma} < \tilde{\rho} = \tilde{\mathcal{O}}((\dot{\varsigma} \tilde{\vee} \dot{\varrho}) \tilde{\vee} \dot{\kappa}) \wedge \tilde{\mathcal{O}}(\dot{\kappa}) \wedge \tilde{\tau}.$$

Then $((\dot{\varsigma} \tilde{\vee} \dot{\varrho}) \tilde{\vee} \dot{\kappa})_{\tilde{\rho}} \in_{\tilde{\sigma}} \tilde{\mathcal{O}}$, $\dot{\kappa}_{\tilde{\rho}} \in_{\sigma} \tilde{\mathcal{O}}$, but $(\dot{\varsigma} \tilde{\vee} (\dot{\varrho} \tilde{\vee} (\dot{\varrho} \tilde{\vee} \dot{\varsigma})))_{\tilde{\rho}} \in_{\tilde{\sigma}} \overline{\vee q_{\tilde{\tau}}} \tilde{\mathcal{O}}$.

Since $\tilde{\mathcal{O}}_{\tilde{\rho}}^{\tilde{\sigma}}$ is a \mathcal{FI} of $\tilde{\mathcal{N}}$, we have $\dot{\varsigma} \tilde{\vee} (\dot{\varrho} \tilde{\vee} (\dot{\varrho} \tilde{\vee} \dot{\varsigma})) \in \tilde{\mathcal{O}}_{\tilde{\rho}}^{\tilde{\sigma}}$, a contradiction.

Hence

$$\tilde{\mathcal{O}}(\dot{\varsigma} \tilde{\vee} (\dot{\varrho} \tilde{\vee} (\dot{\varrho} \tilde{\vee} \dot{\varsigma}))) \vee \tilde{\sigma} \geq \tilde{\mathcal{O}}((\dot{\varsigma} \tilde{\vee} \dot{\varrho}) \tilde{\vee} \dot{\kappa}) \wedge \tilde{\mathcal{O}}(\dot{\kappa}) \wedge \tilde{\tau}.$$

Therefore $\tilde{\delta}$ be a $q\mathcal{P}-(\in_{\tilde{\sigma}}, \in_{\tilde{\sigma}} \vee q_{\tilde{\tau}})\text{-}\mathcal{FFI}$ of $\tilde{\aleph}$.

(2) The proof follows a similar pattern as in (1), and therefore, we omit it for brevity.

(3) Let $\tilde{\delta}$ is a $q\mathcal{P}-(\in_{\tilde{\sigma}}, \in_{\tilde{\sigma}} \vee q_{\tilde{\tau}})\text{-}\mathcal{FFI}$ of $\tilde{\aleph}$ and $\tilde{\rho} \in (\tilde{\sigma}, \hat{1}]$. Then $\forall \zeta \in [\tilde{\delta}]_{\tilde{\rho}}^{\tilde{\tau}}$,

$$\zeta_{\tilde{\rho}} \in_{\tilde{\sigma}} \vee q_{\tilde{\tau}} \tilde{\delta} \Rightarrow \tilde{\delta}(\zeta) \geq \tilde{\rho} > \tilde{\sigma} \text{ or } \tilde{\delta}(\zeta) > 2\tilde{\tau} - \tilde{\rho} > 2\tilde{\tau} - \hat{1} = \tilde{\sigma}.$$

Since $\tilde{\delta}$ is a $q\mathcal{P}-(\in_{\tilde{\sigma}}, \in_{\tilde{\sigma}} \vee q_{\tilde{\tau}})\text{-}\mathcal{FFI}$ of $\tilde{\aleph}$,

$$\tilde{\delta}(0) \vee \tilde{\sigma} \geq \tilde{\delta}(\zeta) \wedge \tilde{\tau} > \tilde{\sigma} \wedge \tilde{\tau} = \tilde{\sigma},$$

and so

$$\tilde{\delta}(0) \geq \tilde{\sigma} \Rightarrow \tilde{\delta}(0) \geq \tilde{\delta}(\zeta) \wedge \tilde{\tau}.$$

Case 1: Let $\tilde{\rho} \in (\tilde{\sigma}, \tilde{\tau}]$. Then $2\tilde{\tau} - \tilde{\rho} \geq \tilde{\tau} \geq \tilde{\rho}$,

$$\tilde{\delta}(0) \geq \tilde{\delta}(\zeta) \wedge \tilde{\tau} \geq \tilde{\rho} \wedge \tilde{\tau} = \tilde{\rho} \text{ or } \tilde{\delta}(0) \geq \tilde{\delta}(\zeta) \wedge \tilde{\tau} > (2\tilde{\tau} - \tilde{\rho}) \wedge \tilde{\tau} = \tilde{\rho} \wedge \tilde{\tau} = \tilde{\rho}.$$

Thus, $0_{\tilde{\rho}} \in_{\tilde{\sigma}} \tilde{\delta}$.

Case 2: Let $\tilde{\rho} \in (\tilde{\tau}, \hat{1}]$. Then $2\tilde{\tau} - \tilde{\rho} < \tilde{\tau} < \tilde{\rho}$,

$$\tilde{\delta}(0) \geq \tilde{\delta}(\zeta) \wedge \tilde{\tau} = \tilde{\rho} \wedge \tilde{\tau} = \tilde{\tau} > 2\tilde{\tau} - \tilde{\rho} \text{ or } \tilde{\delta}(0) \geq \tilde{\delta}(\zeta) \wedge \tilde{\tau} > (2\tilde{\tau} - \tilde{\rho}) \wedge \tilde{\tau} = 2\tilde{\tau} - \tilde{\rho}.$$

Hence, $0_{\tilde{\rho}q_{\tilde{\tau}}\tilde{\delta}} \Rightarrow 0_{\tilde{\rho}} \in_{\tilde{\sigma}} \vee q_{\tilde{\tau}} \tilde{\delta}$.

Let $(\zeta \check{\vee} \varrho) \check{\vee} \kappa, \kappa \in [\tilde{\delta}]_{\tilde{\rho}}^{\tilde{\tau}}$. Then $((\zeta \check{\vee} \varrho) \check{\vee} \kappa)_{\tilde{\rho}, \kappa_{\tilde{\rho}}} \in_{\tilde{\sigma}} \vee q_{\tilde{\tau}} \tilde{\delta}$,

$$\tilde{\delta}((\zeta \check{\vee} \varrho) \check{\vee} \kappa) \geq \tilde{\rho} > \tilde{\sigma} \text{ or } \tilde{\delta}((\zeta \check{\vee} \varrho) \check{\vee} \kappa) > 2\tau - \tilde{\rho} > 2\tilde{\tau} - \hat{1} = \tilde{\sigma}$$

and

$$\tilde{\delta}(\kappa) \geq \tilde{\rho} > \tilde{\sigma} \text{ or } \geq (2\tilde{\tau} - \tilde{\rho}) \wedge \tilde{\tau} \tilde{\delta}(\kappa) > 2\tilde{\tau} - \tilde{\rho} = 2\tilde{\tau} - \tilde{\rho} > 2\tilde{\tau} - \hat{1} = \tilde{\sigma}.$$

Since $\tilde{\delta}$ is a $q\mathcal{P}-(\in_{\tilde{\sigma}}, \in_{\tilde{\sigma}} \vee q_{\tilde{\tau}})\text{-}\mathcal{FFI}$ of $\tilde{\aleph}$,

$$\tilde{\delta}(\zeta \check{\vee} (\varrho \check{\vee} (\varrho \check{\vee} \zeta))) \vee \tilde{\sigma} \geq \tilde{\delta}((\zeta \check{\vee} \varrho) \check{\vee} \kappa) \wedge \tilde{\delta}(\kappa) \wedge \tilde{\tau} > \tilde{\sigma} \wedge \tilde{\sigma} \wedge \tilde{\tau} > \tilde{\sigma} \wedge \tilde{\tau} = \tilde{\sigma}.$$

Therefore,

$$\tilde{\delta}(\zeta \check{\vee} (\varrho \check{\vee} (\varrho \check{\vee} \zeta))) \geq \tilde{\sigma} \Rightarrow \tilde{\delta}(\zeta \check{\vee} (\varrho \check{\vee} (\varrho \check{\vee} \zeta))) \geq \tilde{\delta}((\zeta \check{\vee} \varrho) \check{\vee} \kappa) \wedge \tilde{\delta}(\kappa) \wedge \tilde{\tau}.$$

Case 1: Let $\tilde{\rho} \in (\tilde{\sigma}, \tilde{\tau}]$. Then $2\tilde{\tau} - \tilde{\rho} \geq \tilde{\tau} \geq \tilde{\rho}$,

$$\begin{aligned} \tilde{\delta}(\zeta \check{\vee} (\varrho \check{\vee} (\varrho \check{\vee} \zeta))) &\geq \tilde{\delta}((\zeta \check{\vee} \varrho) \check{\vee} \kappa) \wedge \tilde{\delta}(\kappa) \wedge \tilde{\tau} \\ &\geq \tilde{\rho} \wedge \tilde{\rho} \wedge \tilde{\tau} \\ &= \tilde{\rho} \wedge \tilde{\tau} \\ &= \tilde{\rho} \end{aligned}$$

or

$$\begin{aligned}\tilde{\partial}(\dot{\varsigma} \tilde{\vee} (\dot{\varrho} \tilde{\vee} (\dot{\varrho} \tilde{\vee} \dot{\varsigma}))) &\geq \tilde{\partial}((\dot{\varsigma} \tilde{\vee} \dot{\varrho}) \tilde{\vee} \dot{\kappa}) \wedge \tilde{\partial}(\dot{\kappa}) \wedge \tilde{\tau} \\ &\geq \tilde{\rho} \wedge (2\tilde{\tau} - \tilde{\rho}) \wedge \tilde{\tau} \\ &= \tilde{\rho} \wedge \tilde{\tau} \wedge \tilde{\tau} \\ &= \tilde{\rho} \wedge \tilde{\tau} \\ &= \tilde{\rho}\end{aligned}$$

or

$$\begin{aligned}\tilde{\partial}(\dot{\varsigma} \tilde{\vee} (\dot{\varrho} \tilde{\vee} (\dot{\varrho} \tilde{\vee} \dot{\varsigma}))) &\geq \tilde{\partial}((\dot{\varsigma} \tilde{\vee} \dot{\varrho}) \tilde{\vee} \dot{\kappa}) \wedge \tilde{\partial}(\dot{\kappa}) \wedge \tilde{\tau} \\ &\geq (2\tilde{\tau} - \tilde{\rho}) \wedge (2\tilde{\tau} - \tilde{\rho}) \wedge \tilde{\tau} \\ &= \tilde{\tau} \wedge \tilde{\tau} \wedge \tilde{\tau} \\ &= \tilde{\tau} > \tilde{\rho}.\end{aligned}$$

Hence, $(\dot{\varsigma} \tilde{\vee} (\dot{\varrho} \tilde{\vee} (\dot{\varrho} \tilde{\vee} \dot{\varsigma})))_{\tilde{\rho}} \in_{\tilde{\sigma}} \tilde{\partial}$.

Case 2: Let $\tilde{\rho} \in (\tilde{\tau}, \hat{1}]$. Then $2\tilde{\tau} - \tilde{\rho} < \tilde{\tau} < \tilde{\rho}$,

$$\begin{aligned}\tilde{\partial}(\dot{\varsigma} \tilde{\vee} (\dot{\varrho} \tilde{\vee} (\dot{\varrho} \tilde{\vee} \dot{\varsigma}))) &\geq \tilde{\partial}((\dot{\varsigma} \tilde{\vee} \dot{\varrho}) \tilde{\vee} \dot{\kappa}) \wedge \tilde{\partial}(\dot{\kappa}) \wedge \tilde{\tau} \\ &\geq \tilde{\rho} \wedge \tilde{\rho} \wedge \tilde{\tau} \\ &= \tilde{\rho} \wedge \tilde{\tau} \\ &= \tilde{\tau} > 2\tilde{\tau} - \tilde{\rho}\end{aligned}$$

or

$$\begin{aligned}\tilde{\partial}(\dot{\varsigma} \tilde{\vee} (\dot{\varrho} \tilde{\vee} (\dot{\varrho} \tilde{\vee} \dot{\varsigma}))) &\geq \tilde{\partial}((\dot{\varsigma} \tilde{\vee} \dot{\varrho}) \tilde{\vee} \dot{\kappa}) \wedge \tilde{\partial}(\dot{\kappa}) \wedge \tilde{\tau} \\ &\geq \tilde{\rho} \wedge (2\tilde{\tau} - \tilde{\rho}) \wedge \tilde{\tau} \\ &\geq \tilde{\tau} \wedge (2\tilde{\tau} - \tilde{\rho}) \wedge \tilde{\tau} \\ &\geq \tilde{\tau} \wedge (2\tilde{\tau} - \tilde{\rho}) \\ &= (2\tilde{\tau} - \tilde{\rho})\end{aligned}$$

or

$$\begin{aligned}\tilde{\partial}(\dot{\varsigma} \tilde{\vee} (\dot{\varrho} \tilde{\vee} (\dot{\varrho} \tilde{\vee} \dot{\varsigma}))) &\geq \tilde{\partial}((\dot{\varsigma} \tilde{\vee} \dot{\varrho}) \tilde{\vee} \dot{\kappa}) \wedge \tilde{\partial}(\dot{\kappa}) \wedge \tilde{\tau} \\ &\geq (2\tilde{\tau} - \tilde{\rho}) \wedge (2\tilde{\tau} - \tilde{\rho}) \wedge \tilde{\tau} \\ &\geq (2\tilde{\tau} - \tilde{\rho}) \wedge \tilde{\tau} > (2\tilde{\tau} - \tilde{\rho}).\end{aligned}$$

Thus, $(\dot{\varsigma} \tilde{\vee} (\dot{\varrho} \tilde{\vee} (\dot{\varrho} \tilde{\vee} \dot{\varsigma})))_{\tilde{\rho}q\tilde{\tau}} \tilde{\partial}$.

Hence,

$$(\dot{\varsigma} \tilde{\vee} (\dot{\varrho} \tilde{\vee} (\dot{\varrho} \tilde{\vee} \dot{\varsigma})))_{\tilde{\rho}} \in_{\tilde{\sigma}} \forall q\tilde{\tau} \tilde{\partial} \Rightarrow (\dot{\varsigma} \tilde{\vee} (\dot{\varrho} \tilde{\vee} (\dot{\varrho} \tilde{\vee} \dot{\varsigma}))) \in [\tilde{\partial}]_{\tilde{\rho}}^{\tilde{\tau}}.$$

Therefore $[\tilde{\mathcal{O}}]_{\tilde{\rho}}^{\tilde{\tau}}$ is a \mathcal{FI} of $\tilde{\mathcal{N}}$.

By substituting $\tilde{\sigma} = \tilde{0}$ and $\tilde{\tau} = \widehat{0.5}$ into Theorem 5.4, we can derive the ensuing corollary.

Corollary 6. *Let $\tilde{\mathcal{O}}$ be a $q\mathcal{PF}$ set of $\tilde{\mathcal{N}}$. Then*

- (1) $\tilde{\mathcal{O}}$ is a $q\mathcal{P}-(\in, \in \vee q)\text{-}\mathcal{FFI}$ of $\tilde{\mathcal{N}} \Leftrightarrow \tilde{\mathcal{O}}_{\tilde{\rho}}^{\tilde{\sigma}}(\neq \phi)$ is a \mathcal{FI} of $\tilde{\mathcal{N}}, \forall \tilde{\rho} \in (0, 0.5]^q$.
- (2) $\tilde{\mathcal{O}}$ is a $q\mathcal{P}-(\in, \in \vee q)\text{-}\mathcal{FFI}$ of $\tilde{\mathcal{N}} \Leftrightarrow \langle \tilde{\mathcal{O}} \rangle_{\tilde{\rho}}^{\tilde{\tau}}(\neq \phi)$ is a \mathcal{FI} of $\tilde{\mathcal{N}}, \forall \tilde{\rho} \in (0.5, 1]^q$.
- (3) $\tilde{\mathcal{O}}$ is a $q\mathcal{P}-(\in, \in \vee q)\text{-}\mathcal{FFI}$ of $\tilde{\mathcal{N}} \Leftrightarrow [\tilde{\mathcal{O}}]_{\tilde{\rho}}^{\tilde{\tau}}(\neq \phi)$ is a \mathcal{FI} of $\tilde{\mathcal{N}}, \forall \tilde{\rho} \in (0, 1]^q$.

6. Quadri-Polar Fuzzy TOPSIS Approach

In this section, we present a $q\mathcal{PF}$ TOPSIS approach for multi-criteria group decision-making (MCGDM) problems. For these problems, we use a TOPSIS method based on $q\mathcal{PF}$ -sets to address a set of alternatives $\mathcal{A} = \{\zeta_1, \zeta_2, \zeta_3, \zeta_4\}$ and a set $\mathcal{C} = \{c_1, c_2, c_3, c_4\}$ classified by q . Decision-makers must evaluate the four possibilities based on the $q\mathcal{PF}$ criteria. The possible ratings of alternatives are evaluated in terms of q different attributes among four membership values, represented as $(i = 1, 2, 3, 4)$.

Step 1: The degree of each alternative $\zeta_j \in \mathcal{A}, j = 1, 2, 3, 4$ over all the criteria $(c_k \in \mathcal{C}, k = 1, 2, 3, 4)$ may be expressed as $q\mathcal{PFES}$.

$$\tilde{\mathcal{O}}^{jk}(\zeta) = (\rho_1 \circ \tilde{\mathcal{O}}^{jk}(\zeta), \rho_2 \circ \tilde{\mathcal{O}}^{jk}(\zeta), \rho_3 \circ \tilde{\mathcal{O}}^{jk}(\zeta), \rho_4 \circ \tilde{\mathcal{O}}^{jk}(\zeta)),$$

where $\circ = (\rho_1 \circ \tilde{\mathcal{O}}^{jk}(\zeta) \mid i = 1, 2, \dots, q)$. The tabular representation of the $q\mathcal{PF}$ decision matrix is given by Table 2, which describes the ratings of alternatives.

Table 3. Tabular representation of $q\mathcal{PF}$ decision matrix.

Alternatives	c_1	c_2	c_3	c_4
ζ_1	$\tilde{\mathcal{O}}^{11}(\zeta_1)$	$\tilde{\mathcal{O}}^{12}(\zeta_1)$	$\tilde{\mathcal{O}}^{13}(\zeta_1)$	$\tilde{\mathcal{O}}^{14}(\zeta_1)$
ζ_2	$\tilde{\mathcal{O}}^{21}(\zeta_2)$	$\tilde{\mathcal{O}}^{22}(\zeta_2)$	$\tilde{\mathcal{O}}^{23}(\zeta_2)$	$\tilde{\mathcal{O}}^{24}(\zeta_2)$
ζ_3	$\tilde{\mathcal{O}}^{31}(\zeta_3)$	$\tilde{\mathcal{O}}^{32}(\zeta_3)$	$\tilde{\mathcal{O}}^{33}(\zeta_3)$	$\tilde{\mathcal{O}}^{34}(\zeta_3)$
ζ_4	$\tilde{\mathcal{O}}^{41}(\zeta_4)$	$\tilde{\mathcal{O}}^{42}(\zeta_4)$	$\tilde{\mathcal{O}}^{43}(\zeta_4)$	$\tilde{\mathcal{O}}^{44}(\zeta_4)$

Step 2: We build the optimistic or pessimistic $q\mathcal{PF}$ decision matrix by adding the maximal and smallest values to equalize the length of all $q\mathcal{PFES}$.

Step 3: Weights can be assigned to each $q\mathcal{PF}$ criteria of alternatives by decision-makers based on their choice and importance of each criterion. We assume that the weights assigned by the decision-makers are

$$\mathcal{W} = (w_1, w_2, w_3, w_4) \in (0, 1],$$

satisfying the normalized condition

$$\sum_{k=1}^q w_k = 1, q = 1, 2, 3, 4..$$

Step 4: The weighted $q\mathcal{P}\mathcal{F}$ decision matrix is calculated in Table 4.

Table 4. Tabular representation of a weighted $q\mathcal{P}\mathcal{F}$ decision matrix.

Alternatives	c_1	c_2	c_3	c_4
$\dot{\varsigma}_1$	$\tilde{\partial}(\dot{\varsigma}_1)^{11'}$	$\tilde{\partial}(\dot{\varsigma}_1)^{12'}$	$\tilde{\partial}(\dot{\varsigma}_1)^{13'}$	$\tilde{\partial}(\dot{\varsigma}_1)^{14'}$
$\dot{\varsigma}_2$	$\tilde{\partial}(\dot{\varsigma}_2)^{21'}$	$\tilde{\partial}(\dot{\varsigma}_2)^{22'}$	$\tilde{\partial}(\dot{\varsigma}_2)^{23'}$	$\tilde{\partial}(\dot{\varsigma}_2)^{24'}$
$\dot{\varsigma}_3$	$\tilde{\partial}(\dot{\varsigma}_3)^{31'}$	$\tilde{\partial}(\dot{\varsigma}_3)^{32'}$	$\tilde{\partial}(\dot{\varsigma}_3)^{33'}$	$\tilde{\partial}(\dot{\varsigma}_3)^{34'}$
$\dot{\varsigma}_4$	$\tilde{\partial}(\dot{\varsigma}_4)^{41'}$	$\tilde{\partial}(\dot{\varsigma}_4)^{42'}$	$\tilde{\partial}(\dot{\varsigma}_4)^{43'}$	$\tilde{\partial}(\dot{\varsigma}_4)^{44'}$

For each possible j and k ,

$$\tilde{\partial}^{jk'}(\dot{\varsigma}) = (\rho_1 \circ \tilde{\partial}^{jk'}(\dot{\varsigma}), \rho_2 \circ \tilde{\partial}^{jk'}(\dot{\varsigma}), \rho_3 \circ \tilde{\partial}^{jk'}(\dot{\varsigma}), \rho_4 \circ \tilde{\partial}^{jk'}(\dot{\varsigma})),$$

Step 5: The $q\mathcal{P}\mathcal{F}$ positive ideal solution ($q\mathcal{P}\mathcal{F}PIS$) and $q\mathcal{P}\mathcal{F}$ negative ideal solution ($q\mathcal{P}\mathcal{F}NIS$) of alternatives under the $q\mathcal{P}\mathcal{F}$ environment can be calculated by Equations (5) and (6) as

$$q\mathcal{P} - \mathcal{F}PIS = \{(\tilde{\partial}^1(\dot{\varsigma}))^+, (\tilde{\partial}^2(\dot{\varsigma}))^+, (\tilde{\partial}^3(\dot{\varsigma}))^+, (\tilde{\partial}^4(\dot{\varsigma}))^+\}, \quad (5)$$

$$q\mathcal{P} - \mathcal{F}NIS = \{(\tilde{\partial}^1(\dot{\varsigma}))^-, (\tilde{\partial}^2(\dot{\varsigma}))^-, (\tilde{\partial}^3(\dot{\varsigma}))^-, (\tilde{\partial}^4(\dot{\varsigma}))^-\}, \quad (6)$$

where

$$\begin{aligned} (\tilde{\partial}^{k'}(\dot{\varsigma}))^+ &= \sup_j (\tilde{\partial}^{jk'}(\dot{\varsigma})) \\ &= \sup_j (\rho_1 \circ \tilde{\partial}^{jk'}(\dot{\varsigma}), \rho_2 \circ \tilde{\partial}^{jk'}(\dot{\varsigma}), \rho_3 \circ \tilde{\partial}^{jk'}(\dot{\varsigma}), \rho_4 \circ \tilde{\partial}^{jk'}(\dot{\varsigma})) \\ &= ((\rho_1 \circ \tilde{\partial}^{k'}(\dot{\varsigma}))^+, (\rho_2 \circ \tilde{\partial}^{k'}(\dot{\varsigma}))^+, (\rho_3 \circ \tilde{\partial}^{k'}(\dot{\varsigma}))^+, (\rho_4 \circ \tilde{\partial}^{k'}(\dot{\varsigma}))^+), \end{aligned}$$

and

$$\begin{aligned} (\tilde{\partial}^{k'}(\dot{\varsigma}))^- &= \inf_j (\tilde{\partial}^{jk'}(\dot{\varsigma})) \\ &= \inf_j (\rho_1 \circ \tilde{\partial}^{jk'}(\dot{\varsigma}), \rho_2 \circ \tilde{\partial}^{jk'}(\dot{\varsigma}), \rho_3 \circ \tilde{\partial}^{jk'}(\dot{\varsigma}), \rho_4 \circ \tilde{\partial}^{jk'}(\dot{\varsigma})) \\ &= ((\rho_1 \circ \tilde{\partial}^{k'}(\dot{\varsigma}))^-, (\rho_2 \circ \tilde{\partial}^{k'}(\dot{\varsigma}))^-, (\rho_3 \circ \tilde{\partial}^{k'}(\dot{\varsigma}))^-, (\rho_4 \circ \tilde{\partial}^{k'}(\dot{\varsigma}))^-). \end{aligned}$$

Step 6: The $q\mathcal{P}\mathcal{F}$ Euclidean distance of each alternative $\dot{\varsigma}_j$ from $q\mathcal{P}\mathcal{F}PIS$ and $q\mathcal{P}\mathcal{F}NIS$ can be calculated by Equations (7) and (8).

$$D'_E(\dot{\varsigma}_j, q\mathcal{P} - \mathcal{F}PIS) = \sqrt{\frac{1}{16} \sum_{k=1}^4 [\sum_{l=1}^4 \{ \sum_{i=1}^4 (\rho_i \circ \tilde{\partial}^{jk'}(\dot{\varsigma}) - \rho_i \circ \tilde{\partial}^{k'}(\dot{\varsigma}))^+ \}]}, \quad (7)$$

and

$$D'_E(\zeta_j, q\mathcal{P} - \mathcal{FNIS}) = \sqrt{\frac{1}{16} \sum_{k=1}^4 \left[\sum_{l=1}^4 \left\{ \sum_{i=1}^4 (\rho_i \circ \tilde{\mathfrak{D}}^{jk'}(\zeta) - \rho_i \circ \tilde{\mathfrak{D}}^{k'}(\zeta)) \right\} \right]}, \quad (8)$$

Step 7: The relative $q\mathcal{P}$ - \mathcal{F} closeness coefficient of each alternative ζ_j using the following formula as described by (9),

$$E'_j = \frac{D'_E(\zeta_j, q\mathcal{P} - \mathcal{FNIS})}{D'_E(\zeta_j, q\mathcal{P} - \mathcal{FPIS}) + D'_E(\zeta_j, q\mathcal{P} - \mathcal{FNIS})}, j = 1, 2, 3, 4. \quad (9)$$

The alternative with the highest $q\mathcal{P}$ - \mathcal{F} closeness coefficient is the best one, and we can rank each alternative in order.

We present our proposed decision-making method in Algorithm 1.

In Section 6.1, we examine the practical usage of our suggested model. Specifically, we

Algorithm 1. The algorithm of the proposed approaches for dealing MCGDM problems.

Step 1. Input.

Step 2. Determine the optimistic or pessimistic $q\mathcal{P}\mathcal{F}$ decision matrix.

Step 3. Calculate the normalized weights.

Step 4. Calculate the weight for the pessimistic $q\mathcal{P}$ - \mathcal{F} decision matrix.

Step 5. Compute the $q\mathcal{P}$ - \mathcal{FPIS} , and $q\mathcal{P}$ - \mathcal{FNIS} .

Step 6. $q\mathcal{P}$ - \mathcal{F} Euclidean distance of each alternative ζ_j from $q\mathcal{FPIS}$ and $q\mathcal{P}$ - \mathcal{FPIS} .

Step 7. Calculate the relative $q\mathcal{P}$ - \mathcal{F} closeness coefficients $E'_{j'}$.

Step 8. Output.

Rank the possibilities for the final decision and choose the best one.

show how $q\mathcal{P}$ - \mathcal{F} is useful in the selection of solar power plant stations in a rural region.

Selection of Solar Power Plant Station in a Rural Area

Assume that the government want to build a solar power plant station in a rural area. The government has four options for the location of a new solar power plant. Each site was appraised by a group of decision makers based on the following criteria as follows:

1. "Solar Irradiance (T_1)", which could have the characteristics shown below:
 - Solar Irradiance Density: The amount of solar power received per unit area.
 - Sun Light Duration: The number of sunlight hours per day.
 - Solar Tracking: The technology used to follow the sun's path to maximize energy capture.
 - Temporal Variability: The fluctuation of solar irradiance over time, including daily and seasonal changes.

2. “Land Availability (T_2)”, which could have the characteristics shown below:
 - Land Size: The total area available for the solar power plant.
 - Land Ownership: The ownership status of the land, such as government-owned, privately-owned, or leased.
 - Land Access: The ease of access to the land for construction and maintenance.
 - Land Cost: The cost of acquiring or leasing the land for the project.
3. “Environmental Impacts (T_3)”, which could have the characteristics shown below:
 - Land Use: The current and previous use of the land and the impact of converting it to a solar power plant.
 - Water Consumption: The amount of water required for cleaning solar panels and other operations.
 - Biodiversity: The effect of the solar power plant on local wildlife and plant species.
 - Materials and Waste: The environmental impact of materials used in the construction and the waste generated.
4. “Proximity to Grid Connections (T_4)”, which could have the characteristics shown below:
 - Grid Connection Cost: The expense associated with connecting the solar power plant to the nearest grid infrastructure.
 - Electricity Demand: The local demand for electricity and how the new plant will meet or exceed this demand.
 - Grid Stability: The ability of the existing grid to handle the additional load from the solar power plant.
 - Grid Capacity: The existing capacity of the grid to integrate the new power supply without significant upgrades.
5. “Local Regulations (T_5)”, which could have the characteristics shown below:
 - Zoning Laws: Regulations that dictate land use in the area.
 - Permitting Process: The complexity and duration of obtaining the necessary permits for construction and operation.
 - Incentives and Subsidies: Availability of government incentives, tax credits, and subsidies for renewable energy projects.
 - Compliance Requirements: Environmental and operational regulations that need to be met for the project to proceed.

The five criteria and their attributes are shown below:

1. The $q\mathcal{P}\text{-}\mathcal{F}$ initial decision matrix is represented in Table 5.

Table 5. $q\mathcal{P}\text{-}\mathcal{F}$ initial decision matrix.

Alternatives	$q\mathcal{P}\text{-}\mathcal{F}$ criteria c_1 -Solar Radiance
$\dot{\varsigma}_1$	$\{(.18,.73,.43,.67),(.32,.64,.49,.72),(.38,.66,.54,.64)\}$
$\dot{\varsigma}_2$	$\{(.46,.76,.45,.27),(.56,.91,.36,.48)\}$
$\dot{\varsigma}_3$	$\{(.85,.37,.45,.59),(.72,.48,.72,.58),(.51,.64,.55,.32)\}$
$\dot{\varsigma}_4$	$\{(.21,.52,.34,.77),(.41,.61,.43,.78),(.42,.66,.39,.87)\}$
$\dot{\varsigma}_5$	$\{(.11,.33,.56,.61),(.31,.41,.6,.73)\}$

Alternatives	$q\mathcal{P}\text{-}\mathcal{F}$ criteria c_2 -Local Availability
$\dot{\varsigma}_1$	$\{(.75,.45,.67,.69),(.79,.37,.57,.69)\}$
$\dot{\varsigma}_2$	$\{(.45,.70,.49,.86),(.56,.72,.66,.74),(.47,.62,.58,.72)\}$
$\dot{\varsigma}_3$	$\{(.46,.66,.71,.17),(.41,.77,.78,.19),(.48,.80,.83,.15)\}$
$\dot{\varsigma}_4$	$\{(.57,.64,.38,.57),(.51,.56,.59,.47)\}$
$\dot{\varsigma}_5$	$\{(.24,.12,.81,.77),(.31,.25,.86,.73),(.23,.31,.80,.75)\}$

Alternatives	$q\mathcal{P}\text{-}\mathcal{F}$ criteria c_3 -Environmental Impacts
$\dot{\varsigma}_1$	$\{(.82,.51,.67,.7),(.8,.64,.69,.8),(.7,.59,.68,.78)\}$
$\dot{\varsigma}_2$	$\{(.70,.56,.45,.29),(.73,.51,.37,.43),(.83,.63,.65,.54)\}$
$\dot{\varsigma}_3$	$\{(.74,.46,.61,.8),(.81,.52,.59,.91)\}$
$\dot{\varsigma}_4$	$\{(.66,.55,.46,.88),(.61,.58,.41,.86),(.71,.61,.48,.93)\}$
$\dot{\varsigma}_5$	$\{(.11,.76,.42,.61),(.31,.79,.46,.72),(.23,.8,.57,.84)\}$

Alternatives	$q\mathcal{P}\text{-}\mathcal{F}$ criteria c_4 -Local regulations
$\dot{\varsigma}_1$	$\{(.72,.91,.67,.75),(.69,.84,.69,.87),(.73,.87,.60,.81)\}$
$\dot{\varsigma}_2$	$\{(.60,.44,.76,.80),(.56,.51,.75,.77),(.63,.41,.68,.85)\}$
$\dot{\varsigma}_3$	$\{(.74,.56,.47,.80),(.81,.56,.53,.91),(.61,.60,.65,.89)\}$
$\dot{\varsigma}_4$	$\{(.43,.66,.77,.41),(.51,.68,.76,.36),(.53,.60,.87,.27)\}$
$\dot{\varsigma}_5$	$\{(.21,.65,.52,.61),(.27,.67,.41,.77)\}$

2. The pessimistic decision matrix $q\mathcal{P}\text{-}\mathcal{F}$ in Table 6.

3. The criteria's normalised weights are shown below,

$$w_1 = .2412, w_2 = .2424, w_3 = .2548, w_4 = .2616,$$

where $\sum_{i=1}^4 w_i = 1$.

4. The weight pessimistic of the $q\mathcal{P}\text{-}\mathcal{F}$ decision matrix is calculated in Table 7.

Table 6. The pessimistic $q\mathcal{P}\text{-}\mathcal{F}$ decision matrix.

Alternatives	$q\mathcal{P}\text{-}\mathcal{F}$ criteria c_1 -Solar Radiance
$\dot{\varsigma}_1$	$\{(.18,.73,.43,.67),(.32,.64,.49,.72),(.38,.66,.54,.64)\}$
$\dot{\varsigma}_2$	$\{(.46,.76,.45,.27),(.46,.76,.45,.27),(.56,.91,.36,.48)\}$
$\dot{\varsigma}_3$	$\{(.85,.37,.45,.59),(.72,.48,.72,.58),(.51,.64,.55,.32)\}$
$\dot{\varsigma}_4$	$\{(.21,.52,.34,.77),(.41,.61,.43,.78),(.42,.66,.39,.87)\}$
$\dot{\varsigma}_5$	$\{(.11,.33,.56,.61),(.11,.33,.56,.61),(.31,.41,.6,.73)\}$

Alternatives	$q\mathcal{P}\text{-}\mathcal{F}$ criteria c_2 -Local Availability
$\dot{\varsigma}_1$	$\{(.75,.45,.67,.69),(.79,.37,.57,.69),(.79,.37,.57,.69)\}$
$\dot{\varsigma}_2$	$\{(.45,.7,.49,.86),(.56,.72,.66,.74),(.47,.62,.58,.72)\}$
$\dot{\varsigma}_3$	$\{(.46,.66,.71,.17),(.41,.77,.78,.19),(.48,.80,.83,.15)\}$
$\dot{\varsigma}_4$	$\{(.57,.64,.38,.57),(.51,.56,.59,.47),(.51,.56,.59,.47)\}$
$\dot{\varsigma}_5$	$\{(.24,.12,.81,.77),(.31,.25,.86,.73),(.23,.31,.8,.75)\}$

Alternatives	$q\mathcal{P}\text{-}\mathcal{F}$ criteria c_3 -Environmental Impacts
$\dot{\varsigma}_1$	$\{(.82,.51,.67,.7),(.8,.64,.69,.8),(.7,.59,.68,.78)\}$
$\dot{\varsigma}_2$	$\{(.7,.56,.45,.29),(.73,.51,.37,.43),(.83,.63,.65,.54)\}$
$\dot{\varsigma}_3$	$\{(.74,.46,.61,.8),(.74,.46,.61,.8),(.81,.52,.59,.91)\}$
$\dot{\varsigma}_4$	$\{(.66,.55,.46,.88),(.61,.58,.41,.86),(.71,.61,.48,.93)\}$
$\dot{\varsigma}_5$	$\{(.11,.76,.42,.61),(.31,.79,.46,.72),(.23,.8,.57,.84)\}$

Alternatives	$q\mathcal{P}\text{-}\mathcal{F}$ criteria c_4 -Local regulations
$\dot{\varsigma}_1$	$\{(.72,.91,.67,.75),(.69,.84,.69,.87),(.73,.87,.6,.81)\}$
$\dot{\varsigma}_2$	$\{(.60,.44,.76,.80),(.56,.51,.75,.77),(.63,.41,.68,.85)\}$
$\dot{\varsigma}_3$	$\{(.74,.56,.47,.80),(.81,.56,.53,.91),(.61,.60,.65,.89)\}$
$\dot{\varsigma}_4$	$\{(.43,.66,.77,.41),(.51,.68,.76,.36),(.53,.60,.87,.27)\}$
$\dot{\varsigma}_5$	$\{(.21,.65,.52,.61),(.21,.65,.54,.61),(.27,.67,.41,.77)\}$

5. The evaluation of the $q\mathcal{P}\text{-}\mathcal{F}$ positive ideal solution ($q\mathcal{P}\text{-}\mathcal{F}PIS$) and $q\mathcal{P}\text{-}\mathcal{F}$ negative ideal solution ($q\mathcal{P}\text{-}\mathcal{F}NIS$) is as follows:

$q\mathcal{P}\text{-}\mathcal{F}PIS =$

$$\begin{aligned} & \{ \{ (.2050, .1833, .1351, .1857), (.1737, .1833, .1737, .1881), (.1351, .2195, .1447, .2098) \}, \\ & \{ (.1818, .1697, .1963, .2085), (.1915, .1866, .2085, .1794), (.1915, .1939, .2012, .1818) \}, \\ & \{ (.2089, .1936, .1707, .2242), (.2038, .2013, .1758, .2191), (.2115, .2038, .1733, .2370) \}, \\ & \{ (.1936, .2381, .2014, .2093), (.2119, .2197, .1988, .2381), (.1910, .2276, .2276, .2328) \} \}, \end{aligned}$$

and

Table 7. The weighted pessimistic $q\mathcal{P}\text{-}\mathcal{F}$ decision matrix.

ζ	$q\mathcal{P}\text{-}\mathcal{F}$ criteria c_1 -Solar Radiance
ζ_1	$\{(.0434,.1761,.1037,.1568),(.0724,.1544,.1182,.1688),(.0917,.1592,.1302,.1544)\}$
ζ_2	$\{(.1110,.1833,.1085,.0651),(.1110,.1833,.1085,.0651),(.1351,.2195,.0868,.1158)\}$
ζ_3	$\{(.2050,.0892,.1085,.1375),(.1737,.1158,.1737,.1399),(.1230,.1544,.1327,.0772)\}$
ζ_4	$\{(.0507,.1254,.0820,.1857),(.0989,.1471,.1037,.1881),(.1013,.1592,.0941,.2098)\}$
ζ_5	$\{(.0265,.0796,.1351,.1471),(.0265,.0796,.1351,.1471),(.0748,.0989,.1447,.1761)\}$
ζ	$q\mathcal{P}\text{-}\mathcal{F}$ criteria c_2 -Local Availability
ζ_1	$\{(.1818,.1091,.1624,.1673),(.1915,.0897,.1382,.1673),(.1915,.0897,.1382,.1673)\}$
ζ_2	$\{(.1091,.1697,.1188,.2085),(.1357,.1745,.1600,.1794),(.1139,.1503,.1406,.1745)\}$
ζ_3	$\{(.1115,.1600,.1721,.0412),(.0994,.1866,.1891,.0461),(.1164,.1939,.2012,.0364)\}$
ζ_4	$\{(.1382,.1551,.0921,.1382),(.1236,.1357,.1430,.1139),(.1236,.1357,.1430,.1139)\}$
ζ_5	$\{(.0582,.0291,.1963,.1866),(.0751,.0606,.2085,.1770),(.0558,.0751,.1939,.1818)\}$
ζ	$q\mathcal{P}\text{-}\mathcal{F}$ criteria c_3 -Environmental Impacts
ζ_1	$\{(.2089,.1299,.1707,.1784),(.2038,.1631,.1758,.2038),(.1784,.1503,.1733,.1987)\}$
ζ_2	$\{(.1784,.1427,.1147,.0739),(.1860,.1299,.0943,.1096),(.2115,.1605,.1656,.1376)\}$
ζ_3	$\{(.1886,.1172,.1554,.2038),(.1886,.1172,.1554,.2038),(.2064,.1325,.1503,.2319)\}$
ζ_4	$\{(.1682,.1401,.1172,.2242),(.1554,.1478,.1045,.2191),(.1809,.1554,.1223,.2370)\}$
ζ_5	$\{(.0280,.1936,.1070,.1554),(.0790,.2013,.1172,.1835),(.0586,.2038,.1452,.2140)\}$
ζ	$q\mathcal{P}\text{-}\mathcal{F}$ criteria c_4 -Local regulations
ζ_1	$\{(.1884,.2381,.1753,.1962),(.1805,.2197,.1805,.2276),(.1910,.2276,.1570,.2119)\}$
ζ_2	$\{(.1570,.1151,.1988,.2093),(.1465,.1334,.1962,.2014),(.1648,.1073,.1779,.2224)\}$
ζ_3	$\{(.1936,.1465,.1230,.2093),(.2119,.1465,.1386,.2381),(.1596,.1570,.1700,.2328)\}$
ζ_4	$\{(.1125,.1727,.2014,.1073),(.1334,.1779,.1988,.0942),(.1386,.1570,.2276,.0706)\}$
ζ_5	$\{(.0549,.1700,.1360,.1596),(.0549,.1700,.1413,.1596),(.0706,.1753,.1073,.2014)\}$

$q\mathcal{P}\text{-}\mathcal{FNIS} =$

$$\begin{aligned} & \{ \{ (.0265, .0796, .0820, .0651), (.0265, .0796, .1037, .0651), (.0748, .0989, .0868, .0772) \}, \\ & \{ (.0582, .0291, .0921, .0412), (.0751, .0606, .1430, .0461), (.0558, .0751, .1406, .0364) \}, \\ & \{ (.0280, .1172, .1070, .0739), (.0790, .1172, .0943, .1096), (.0586, .1325, .1223, .1376) \}, \\ & \{ (.0549, .1151, .1230, .1073), (.0549, .1334, .1386, .0942), (.0706, .1073, .1073, .0706) \} \}, \end{aligned}$$

6. Using (7) and (8), the $q\mathcal{P}\text{-}\mathcal{F}$ Euclidean distance of each alternative ζ_j from $q\mathcal{FPIS}$ and $q\mathcal{P}\text{-}\mathcal{FPIS}$ are calculated as:

$$\begin{aligned} D'_E(\zeta_1, q\mathcal{P}\text{-}\mathcal{FPIS}) &= .0951, \\ D_E(\zeta_2, q\mathcal{P}\text{-}\mathcal{FPIS}) &= .1293, \\ D_E(\zeta_3, q\mathcal{P}\text{-}\mathcal{FPIS}) &= .1229, \\ D_E(\zeta_4, q\mathcal{P}\text{-}\mathcal{FPIS}) &= .1311, \end{aligned}$$

$$D'_E(\zeta_5, q\mathcal{P}\text{-}\mathcal{FPLS}) = .1786,$$

and

$$\begin{aligned} D'_E(\zeta_1, q\mathcal{P}\text{-}\mathcal{FNIS}) &= .1817, \\ D'_E(\zeta_2, q\mathcal{P}\text{-}\mathcal{FNIS}) &= .1627, \\ D'_E(\zeta_3, q\mathcal{P}\text{-}\mathcal{FNIS}) &= .1740, \\ D'_E(\zeta_4, q\mathcal{P}\text{-}\mathcal{FNIS}) &= .1461, \\ D'_E(\zeta_5, q\mathcal{P}\text{-}\mathcal{FNIS}) &= .1218. \end{aligned}$$

Using Equation (9), the relative $q\mathcal{P}\text{-}\mathcal{F}$ closeness coefficients $E_{j'}$ are calculated as:

$$\begin{aligned} E_{1'} &= .6565, \\ E_{2'} &= .5572, \\ E_{3'} &= .5860, \\ E_{4'} &= .5270, \\ E_{5'} &= .4055. \end{aligned}$$

According to the foregoing computations, the final ranking of power plant selection is as follows:

$$\zeta_1 > \zeta_3 > \zeta_2 > \zeta_4 > \zeta_5.$$

Hence, solar power plant station ζ_1 is selected in the rural area.

As a result of the evaluation among the alternatives, the most risky structure is ζ_1 . The positive ideal solution, negative ideal solution and ranking of alternatives based on closeness coefficients is shown in Figure 1.




Fig. 1: Positive ideal solution, Negative ideal solution and Ranking of alternatives

7. Conclusions

Quadri-polar structures are often preferred over clear-cut circumstances. Various forms of information can be used to manipulate the membership degrees to facilitate the handling of quadri-polar fuzzy information. We have proposed a novel class of generalized $q\mathcal{P}$ - $\mathcal{FFI}(s)$ of $\tilde{\mathfrak{N}}$ called, a $q\mathcal{P}$ -(ϖ, ϑ)- $\mathcal{FFI}(s)$ and compared to the generalizations of present fuzzy sets, it is shown to be a more flexible approach that can be evaluated in quadri-polar ways based on practical interests and requirements. We defined and analyzed the concept of $q\mathcal{P}$ -($\in_{\tilde{\sigma}}, \in_{\tilde{\sigma}} \vee q_{\tilde{\tau}}$)- $\mathcal{FFI}(s)$. Moreover, we presented various characterizations of $q\mathcal{P}$ -($\in_{\tilde{\sigma}}, \in_{\tilde{\sigma}} \vee q_{\tilde{\tau}}$)- $\mathcal{FFI}(s)$. It is used to manage data that includes quadri-polar information suggested by decision-makers. The final decision on the proposed approach is determined by the decision-maker's optimistic or pessimistic outlook. In practical perspective, we have devised a q - \mathcal{PF} TOPSIS approach to address MCGDM problems. This method represents a natural extension of the TOPSIS method tailored to our specific model. Ultimately, we have implemented our methodology in addressing real-world issues. In the future, we will delve into additional decision-making methods associated with the proposed concept, such as q -polar fuzzy semi-hyper groups, q -polar fuzzy rough sets, q -polar fuzzy in different logical algebras environment.

Acknowledgements

The authors gratefully acknowledge the funding of the Deanship of Graduate Studies and Scientific Research, Jazan University, Saudi Arabia, through Project Number: GSSRD-24.

References

- [1] Saeid A.B. Fantastic ideals in bci-algebras. *World Applied Sciences Journal*, 8(5):550–554, 2010.
- [2] E.A. Abuhijleh, M. Massadeh, A. Sheimat, and A. Alkouri. Complex fuzzy groups based on rosenfled's approach. *WSEAS Trans. Math.*, 20:368–377, 2021.
- [3] M. Akram and A. Adeel. Novel topsis method for group decision-making based on hesitant m-polar fuzzy model. *J. Int. Fuzzy syst.*, 37(6):8077–8096, 2019.
- [4] M. Akram, A. Farooq, and K. P. Shum. On m -polar fuzzy lie subalgebras. *Italian J. Pure Appl. Math.*, 36:445–454, 2016.
- [5] A. Al-Masarwah and Ahmad A.G. On some properties of doubt bipolar fuzzy h-ideals in bck/bci-algebras. *Eur. J. Pure Appl. Math.*, 11(3):652–670, 2018.
- [6] A. Al-Masarwah and A.G. Ahmad. m -polar (α, β) -fuzzy ideals in bck/bci-algebras. *Symmetry*, 11(1):1–18, 2019.

- [7] A. Al-Masarwah and A.G. Ahmad. m -polar fuzzy ideals of bck/bci-algebras. *J. King Saud Univ. Sci.*, 31(4):1220–1226, 2019.
- [8] A. Al-Masarwah and A.G. Ahmad. Subalgebras of type (α, β) based on m -polar fuzzy points in bck/bci-algebras. *AIMS Mathematics*, 5(2):1035–1049, 2020.
- [9] Moin A. Ansari, A. Haider, and A.N.A. Koam. On a graph associated to up-algebras. *Mathematical and Computational Applications*, 23(4):61, 2018.
- [10] M. Balamurugan, N. Alessa, K. Loganathan, and N. Amar Nath. $(\acute{e}, \acute{e} \vee \acute{q}_k)$ -uni-intuitionistic fuzzy soft h -ideals in subtraction bg-algebras. *Mathematics*, 11(10), 2296:1–15, 2023.
- [11] M. Balamurugan, N. Alessa, K. Loganathan, and M. Sudheer Kumar. Bipolar intuitionistic fuzzy soft ideals of bck/bci-algebras and its applications in decision-making. *Mathematics*, 11(21):4471, 2023.
- [12] S.K. Bhakat and P. Das. $(\in, \in \vee q)$ -fuzzy subgroups. *Fuzzy Sets Syst.*, 80(3):359–368, 1996.
- [13] J. Chen, S. Li, S. Ma, and X. Wang. m -polar fuzzy sets an extension of bipolar fuzzy sets. *The Sci World J.*, Article Id 416530:1–8, 2014.
- [14] W.A. Dudek, M. Shabir, and M. Irfan Ali. (α, β) -fuzzy ideals of hemirings. *Comput Math Appl.*, 58(2):310–321, 2009.
- [15] A. Fallath, M.O. Massadeh, and A.U. Alkouri. Normal and cosets of (γ, δ) -fuzzy h x-subgroups. *J. Appl. Math. Inform.*, 40:719–727, 2022.
- [16] A. Farooq, G. Alia, and M. Akram. On m -polar fuzzy groups. *Int. J. Algebra Stat.*, 5(2):115–127, 2016.
- [17] K. Hayat, T. Mahmood, and B.Y. Cao. On bipolar anti fuzzy h -ideals in hemi-rings. *Fuzzy Inf. Eng.*, 9(1):1–19, 2017.
- [18] A. Iamapan, M. Balamurugan, and V. Govindan. $(\in, \in \vee q_k)$ -anti-intuitionistic fuzzy soft b -ideals in bck/bci-algebras. *Mathematics and Statistics*, 10(3):515–522, 2022.
- [19] M. Ibrar, A. Khan, and B. Davvaz. Characterizations of regular ordered semigroups in terms of (α, β) -bipolar fuzzy generalized bi-ideals. *J. Intelli Fuzzy Syst.*, 33:365–376, 2017.
- [20] Y. Imai and K. Iseki. On axiom systems of propositional calculi. i. *Proc. Japan Acad.*, 41(6):436–439, 1965.
- [21] K. Iseki. An algebra related with a propositional calculus. *Proc. Japan Acad.*, 42(1):26–29, 1966.

- [22] K. Iseki. On bci-algebras. *Math. Seminar Notes*, 8(1):125–130, 1980.
- [23] K. Iseki and Tanaka S. An introduction to the theory of bck-algebras. *Math Japan*, 23(1), 1978.
- [24] C. Jana and M. Pal. $(\in_\gamma, \in_\gamma \vee q_\delta)$ -fuzzy soft bci-algebras. *Missouri J. Math Sci.*, 29(2):197–215, 2017.
- [25] C. Jana, M. Pal, and A.B. Saeid. $(\in, \in \vee q)$ -bipolar fuzzy bck/bci-algebras. *Missouri J. Math. Sci.*, 29:1–23, 2017.
- [26] Y.B. Jun. On (α, β) -fuzzy ideals of bck/bci-algebras. *Sci. Math. Jpn.*, pages 101–105, 2004.
- [27] Y.B. Jun. On (α, β) -fuzzy subalgebras of bck/bci-algebras. *Bull Korean Math Soc.*, 42(4):703–711, 2005.
- [28] Y.B. Jun. Fuzzy subalgebras of type (α, β) -fuzzy subalgebras in bck/bci-algebras. *KYUNGPOOK Math. J.*, 4:403–410, 2007.
- [29] Kavikumar, Azme Khamis, and Young Bae Jun. Fuzzy bi-ideals in ternary semirings. *International Journal of Computational and Mathematical Sciences*, 3(4):164 – 168, 2009.
- [30] O. Kazanc, S. Hoskova-Mayerova, and B. Davvaz. Multipolar fuzzy hyperideals in ordered semihypergroups. *Mathematics*, pages 1–11, 2022.
- [31] Zadeh L.A. Fuzzy sets. *Information and Control*, 8(3):338–353, 1965.
- [32] K.J. Lee. Bipolar fuzzy subalgebras and bipolar fuzzy ideals of bck/bci-algebras. *Bull. Malays. Math. Sci. Soc.*, 32(3):361–373, 2009.
- [33] X. Ma, J. Zhan, and Y.B. Jun. Some kinds of $(\in_\gamma, \in_\gamma \vee q_\delta)$ -fuzzy ideals of bci-algebras. *Comput Math Appl.*, 61(4):1005–1015, 2011.
- [34] G. Muhiuddin, N. Abughazalah, A. Aljuhani, and M. Balamurugan. Tripolar picture fuzzy ideals of bck-algebras. *Symmetry*, 14(8), 1562:1–20, 2022.
- [35] G. Muhiuddin and A.M. Al-Roqi. Subalgebras of bck/bci-algebras based on (α, β) -type fuzzy sets. *Comput Anal Appl*, 18(6):1057–1064, 2015.
- [36] G. Muhiuddin, R.A. Takallo, M.M. nd Borzooei, and Y.B. Jun. m -polar fuzzy q -ideals in bci-algebras. *J. King Saud Univ. Sci.*, 32(6):2803–2809, 2020.
- [37] A. L. Narayanan and T. Manikantan. $(\in, \in \vee q)$ -fuzzy subnearrings and $(\in, \in \vee q)$ -fuzzy ideals of near-rings. *J. Appl Math Comput.*, 18(1):419–430, 2009.
- [38] A. Rosenfeld. Fuzzy groups. *J. Math Anal Appl.*, 35(3):512–517, 1971.

- [39] Zhang W.R. Bipolar fuzzy sets and relations a computational framework for cognitive and modeling and multiagent decision analysis. *Proceedings of the First International Joint Conference of The North American Fuzzy Information Processing Society Biannual Conference, The Industry Fuzzy Control and Intelligent*, pages 305–309, 1994.
- [40] O.G. Xi. Fuzzy bck-algebras. *Math. Jpn.*, 36:935–942, 1991.
- [41] J. Zhan. Fuzzy soft γ -hyperrings. *Iran J. Sci Technol.*, 36(2):125–135, 2012.
- [42] M. Zulfiqar. Some characterizations of $(\overline{\in}_\gamma, \overline{\in}_\gamma, \vee q_\delta)$ -fuzzy fantastic ideals in bch-algebras. *Acta Sci Technol.*, 35(1):123–129, 2013.
- [43] M. Zulfiqar and M. Shabir. $(\in_\gamma, \in_\gamma \vee q_\delta)$ -fuzzy soft bci-algebras. *University Politehnica of Bucharest Scientific Bulletin-Series A-Applied Mathematics and Physics*, 75(4):217–230, 2013.