

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/382308888>

Crossing cubic Lie algebras

Article in *AIMS Mathematics* · July 2024

DOI: 10.3934/math.20241075

CITATIONS

0

READS

85

4 authors, including:



Anas Al-Masarwah

Ajloun National University

28 PUBLICATIONS 368 CITATIONS

[SEE PROFILE](#)



Majdoleen Abuqamar

Jadara University

16 PUBLICATIONS 260 CITATIONS

[SEE PROFILE](#)



Kholood Alsager

Qassim University

23 PUBLICATIONS 31 CITATIONS

[SEE PROFILE](#)



Research article**Crossing cubic Lie algebras****Anas Al-Masarwah¹, Nadeen Kdaisat², Majdoleen Abuqamar³ and Kholood Alsager^{4,*}**

¹ Department of Mathematics, Faculty of Science, Ajloun National University, P. O. Box 43, Ajloun 26810, Jordan; anas.almasarwah@anu.edu.jo

² Department of Mathematics, Yarmouk University, Shafiq Irshidat Street, Irbid 21163, Jordan; nadeengdesat1@gmail.com

³ Department of Mathematics, Faculty of Science and Information Technology, Jadara University, Irbid 21110, Jordan; mjabuqamar@gmail.com

⁴ Department of Mathematics, College of Science, Qassim University, Buraydah, Saudi Arabia

* **Correspondence:** Email: ksakr@qu.edu.sa; Tel: +962777985088.

Abstract: An interval-valued fuzziness structure is an effective approach addressing ambiguity and for expressing people's hesitation in everyday situations. An \mathcal{N} -structure is a novel technique for solving practical problems. This is beneficial for resolving a variety of issues, and a lot of progress is being made right now. In order to develop crossing cubic structures (CCSs), Jun et al. amalgamate interval-valued fuzziness and \mathcal{N} -structures. In this manuscript, our main contribution is to originate the concepts of crossing cubic (CC) Lie algebra, CC Lie sub-algebra, ideal, and homomorphism. We investigate some properties of these concepts. In a Lie algebra, the construction of a quotient Lie algebra via the CC Lie ideal is provided. Furthermore, the CC isomorphism theorems are presented.

Keywords: Lie algebras; crossing cubic Lie algebras; crossing cubic fields; crossing cubic solvable Lie algebras; crossing cubic nilpotent Lie algebras; fuzzy logic

Mathematics Subject Classification: 06F35, 03G25, 03B52, 03B05

1. Introduction

Algebraic structures are a crucial aspect of pure and applied mathematics, and other domains, with an enormous variety of applications in computer science [1–3], theoretical physics [4], coding theory [5], and other domains. The idea of Lie algebra was initially discovered by Sophus Lie [6] as continued transformation groups (now referred to as Lie groups). The inception of this idea of Lie algebras was presented when Sophus Lie tried to distinguish certain “smooth” subgroups of general linear groups. Many scholars have focused on Lie algebras over the years, creating a variety of

structural concepts pertaining to the theory of groups, rings, fields, and other algebraic aspects. Lie groups, Lie similarities and Lie algebras are fundamental notions in mathematics.

The inception of the notion of fuzziness structures was performed by a well-known mathematician, Zadeh in 1965 [7]. After him, the concept has undergone plenty of additions, improvements and developments by numerous researchers who adopted further advanced and novel kinds and aspects of the notion, such as interval-valued fuzziness structures [8], cubic structures [9], and \mathcal{N} -structures [10]. For the results on algebraic structures with uncertainty (see works by the authors of [11–17]).

In the fuzzification of Lie algebras and bringing together the concepts of fuzziness structure and Lie subalgebras or ideals, the thought of fuzziness structure was connected with Lie algebras by Yehia [18]. A lot of researchers concentrated on Lie algebras in fuzziness structures (see works by the authors of [19–21]). Applying the idea of interval-valued fuzziness structures to Lie algebras, Akram [22] extended the work of [18] and studied the conception of interval-valued fuzziness Lie subalgebras and Lie ideals. In the intuitionistic fuzziness Lie algebras, Akram and Shum [23] applied intuitionistic fuzziness structures to Lie algebras by merging the notions of intuitionistic fuzziness structures and Lie algebras. They proposed and discussed the idea of intuitionistic fuzzy Lie subalgebras and investigated some of their characteristics and properties in a Lie algebra. As an extension of the results in [18, 23], Akram et al. [24] developed single-valued neutrosophic Lie subalgebras and ideals. They described some significant results of single-valued neutrosophic Lie ideals.

In terms of the case of CCS s, Jun et al. [25] were the first one presenting this concept. They defined the same or opposite direction order, S -intersection (union), and O -intersection (union) of CCS s, and discussed their related properties. On a nonempty set \mathcal{L} , a CCS indicates a map known as the membership function

$$\mathcal{C} : \mathcal{L} \rightarrow \mathbb{P}[0, 1] \times [-1, 0].$$

The introduction of CCS s is based on the ideas of interval-valued fuzziness structure indicates a map known as the membership function

$$\widetilde{\omega}_{\mathcal{C}} : \mathcal{L} \rightarrow \mathbb{P}[0, 1]$$

and the negative version of fuzziness structures (say, \mathcal{N} -structures) indicates a map known as the membership function

$$\pi_{\mathcal{C}} : \mathcal{L} \rightarrow [-1, 0].$$

In [26], Jun and Song applied this notion to BCK and BCI-algebras. They implemented the notions of CC ideals in a BCK/BCI-algebra, CC \circ -subalgebras of a BCK-algebra with the condition (S), and investigated certain characterizations and translations of CC subalgebras and ideals. Öztürk et al. [27] applied this notion to commutative BCK-algebras and semigroups. Mostafa et al. [28] proposed $(\bar{\alpha}, \alpha)$ - CC QS-ideals of QS-algebras, and discussed certain related characteristics and results.

To convey the novelty of this framework, Figure 1 depicts a novel hybrid modification of interval-valued fuzziness Lie algebras and a negative version of fuzziness Lie algebras known as CC Lie algebras.




Figure 1. Contributions toward CC Lie algebras.

This study is the initial effort to discuss and utilize the *CCSs* in Lie algebras. The overall structure of the manuscript is as follows: In Section 2, we review the definitions of Lie subalgebra, Lie ideal, and homomorphism. We present the definitions of fuzziness Lie subalgebra and Lie ideal of Lie algebras. Moreover, we propose the notion of *CCSs* with their level cuts. In Section 3, we develop the concepts of CC Lie subalgebras, CC Lie ideals, and homomorphism. We discuss the Cartesian product of CC Lie subalgebras. Also, we describe certain significant results of CC Lie ideals. In Section 4, we provide a construction of a quotient Lie algebra via the CC Lie ideal in a Lie algebra. Also, the CC isomorphism theorems are presented. Lastly, the conclusions and certain potential future studies of the current article are proposed in Section 5.

2. Preliminaries

Here, we initially review some fundamental aspects that are significant for this manuscript. Throughout this manuscript, \mathcal{L} is a Lie algebra.

A Lie algebra is presented as a pair $(\mathcal{L}, [\cdot, \cdot])$, where \mathcal{L} is a vector space over a field F (\mathbb{R} or \mathbb{C}) and $[\cdot, \cdot] : \mathcal{L} \times \mathcal{L} \rightarrow \mathcal{L}$ is denoted by $(l_1, l_2) \rightarrow [l_1, l_2]$ achieving the axioms below:

(A1) $[l_1, l_2]$ bilinear;

(A2) $[l_1, l_2] = 0 \ \forall l_1 \in \mathcal{L}$;

(A3) $[[l_1, l_2], l_3] + [[l_2, l_3], l_1] + [[l_3, l_1], l_2] = 0 \ \forall l_1, l_2, l_3 \in \mathcal{L}$.

We note that the operation “ \cdot ” in \mathcal{L} is anti-commutative, i.e., $[l_1, l_2] = -[l_2, l_1]$. But it is not associative, i.e., $[[l_1, l_2], l_3] \neq [l_1, [l_2, l_3]]$.

Now, we recall the concepts of a Lie subalgebra and a Lie ideal.

- A Lie subalgebra is a subspace \mathcal{H} of \mathcal{L} that is closed under “[\cdot, \cdot]”.
- A Lie ideal is a subspace \mathcal{I} of \mathcal{L} with the property $[\mathcal{I}, \mathcal{L}] \subseteq \mathcal{I}$.
- Any Lie ideal \mathcal{I} is a Lie subalgebra.

Let ϖ be a fuzzy structure on \mathcal{L} , i.e., $\varpi : \mathcal{L} \rightarrow [0, 1]$. Then, ϖ is a fuzzy Lie subalgebra [18] of \mathcal{L} if the following identities are satisfied: $\forall l_1, l_2 \in \mathcal{L}, \alpha \in F$,

$$(S1) \varpi(l_1 + l_2) \geq \min\{\varpi(l_1), \varpi(l_2)\},$$

$$(S2) \varpi(\alpha l_1) \geq \varpi(l_1),$$

$$(S3) \varpi([l_1, l_2]) \geq \min\{\varpi(l_1), \varpi(l_2)\}.$$

A fuzzy structure ϖ is a fuzzy Lie ideal [18] of \mathcal{L} if (S1), (S2), and $\varpi([l_1, l_2]) \geq \varpi(l_1)$ are satisfied $\forall l_1, l_2 \in \mathcal{L}, \alpha \in F$.

Consider the interval number $\tilde{\xi} = [\xi^-, \xi^+]$ of $[0, 1]$, where $\xi^-, \xi^+ \in [0, 1]$ and $\xi^- \leq \xi^+$. $\mathbb{P}[0, 1]$ denotes the set of all interval numbers. For the interval numbers $\tilde{\xi}_i = [\xi_i^-, \xi_i^+], \tilde{\tau}_i = [\tau_i^-, \tau_i^+] \in \mathbb{P}[0, 1]$, where $i \in \Lambda$. We give:

- $\widetilde{\min}\{\tilde{\xi}_i, \tilde{\tau}_i\} = [\min\{\xi_i^-, \tau_i^-\}, \min\{\xi_i^+, \tau_i^+\}]$,
- $\widetilde{\max}\{\tilde{\xi}_i, \tilde{\tau}_i\} = [\max\{\xi_i^-, \tau_i^-\}, \max\{\xi_i^+, \tau_i^+\}]$,
- $\tilde{\xi}_1 \leq \tilde{\xi}_2 \Leftrightarrow \xi_1^- \leq \xi_2^- \text{ and } \xi_1^+ \leq \xi_2^+$,
- $\xi_1 = \xi_2 \Leftrightarrow \xi_1^- = \xi_2^- \text{ and } \xi_1^+ = \xi_2^+$.

To say that $\tilde{\xi}_1 < \tilde{\xi}_2$ (resp. $\tilde{\xi}_1 > \tilde{\xi}_2$), we mean $\tilde{\xi}_1 \leq \tilde{\xi}_2$ and $\tilde{\xi}_1 \neq \tilde{\xi}_2$ (resp. $\tilde{\xi}_1 \geq \tilde{\xi}_2$ and $\tilde{\xi}_1 \neq \tilde{\xi}_2$).

$\mathcal{F}(\mathcal{L}, [-1, 0])$ indicates the collection of maps from a set $\mathcal{L} \neq \emptyset$ to $[-1, 0]$. An object of $\mathcal{F}(\mathcal{L}, [-1, 0])$ is a negative-valued map from \mathcal{L} to $[-1, 0]$ (simply, an \mathcal{N} -function on \mathcal{L}). (\mathcal{L}, π) is an \mathcal{N} -structure, where π is an \mathcal{N} -function on \mathcal{L} .

Definition 2.1 ([25]). Let \mathcal{L} be a non-empty set. A CCS of \mathcal{L} is a structure having the shape:

$$\mathcal{C} = \{\langle l, \widetilde{\varpi}_{\mathcal{C}}(l), \pi_{\mathcal{C}}(l) \rangle \mid l \in \mathcal{L}\},$$

where $\widetilde{\varpi}_{\mathcal{C}} : \mathcal{L} \rightarrow \mathbb{P}[0, 1]$ is an interval valued fuzziness structure on \mathcal{L} and $\pi_{\mathcal{C}} : \mathcal{L} \rightarrow [-1, 0]$ is an \mathcal{N} -function on \mathcal{L} .

For the sake of simplicity, we shall use the symbol $\mathcal{C} = \langle \widetilde{\varpi}_{\mathcal{C}}, \pi_{\mathcal{C}} \rangle$ for the CCS $\mathcal{C} = \{\langle l, \widetilde{\varpi}_{\mathcal{C}}(l), \pi_{\mathcal{C}}(l) \rangle \mid l \in \mathcal{L}\}$.

Definition 2.2 ([25]). Let $\mathcal{C} = \langle \widetilde{\varpi}_{\mathcal{C}}, \pi_{\mathcal{C}} \rangle$ be a CCS of \mathcal{L} . Then, $(\bar{\nu}, \gamma)$ -level of $\mathcal{C} = \langle \widetilde{\varpi}_{\mathcal{C}}, \pi_{\mathcal{C}} \rangle$ is the crisp set in \mathcal{L} denoted by $\Upsilon(\mathcal{C}; (\bar{\nu}, \gamma))$ and is defined as

$$\Upsilon(\mathcal{C}; (\bar{\nu}, \gamma)) = \{l \in \mathcal{L} \mid \widetilde{\varpi}_{\mathcal{C}}(l) \geq \bar{\nu}, \pi_{\mathcal{C}}(l) \leq \gamma\},$$

where $\bar{\nu} \in \mathbb{P}[0, 1]$ and $\gamma \in [-1, 0]$.

For $\bar{\nu} \in \mathbb{P}[0, 1]$ and $\gamma \in [-1, 0]$. The $\bar{\nu}$ -level $\Upsilon(\widetilde{\varpi}_{\mathcal{C}}; \bar{\nu})$ and γ -level $\Upsilon(\pi_{\mathcal{C}}; \gamma)$ subsets of \mathcal{C} can be defined as:

$$\Upsilon(\widetilde{\varpi}_{\mathcal{C}}; \bar{\nu}) = \{l \in \mathcal{L} \mid \widetilde{\varpi}_{\mathcal{C}}(l) \geq \bar{\nu}\} \text{ and } \Upsilon(\pi_{\mathcal{C}}; \gamma) = \{l \in \mathcal{L} \mid \pi_{\mathcal{C}}(l) \leq \gamma\}.$$

3. Crossing cubic Lie subalgebras and ideals

In the current section, we originate the concepts of CC Lie sub-algebra, ideal, and homomorphism. Also, we investigate some features of these notions.

Definition 3.1. A CCS $\mathcal{C} = \langle \widetilde{\varpi}_{\mathcal{C}}, \pi_{\mathcal{C}} \rangle$ is a CC Lie subalgebra of \mathcal{L} if it meets the following identities:

- (A) $(\forall l_1, l_2 \in \mathcal{L}) \left(\begin{array}{l} \widetilde{\omega}_{\mathcal{C}}(l_1 + l_2) \geq \widetilde{\min}\{\widetilde{\omega}_{\mathcal{C}}(l_1), \widetilde{\omega}_{\mathcal{C}}(l_2)\}, \\ \pi_{\mathcal{C}}(l_1 + l_2) \leq \max\{\pi_{\mathcal{C}}(l_1), \pi_{\mathcal{C}}(l_2)\} \end{array} \right),$
- (B) $(\forall l_1 \in \mathcal{L}, \alpha \in F) \left(\begin{array}{l} \widetilde{\omega}_{\mathcal{C}}(\alpha l_1) \geq \widetilde{\omega}_{\mathcal{C}}(l_1), \\ \pi_{\mathcal{C}}(\alpha l_1) \leq \pi_{\mathcal{C}}(l_1) \end{array} \right),$
- (C) $(\forall l_1, l_2 \in \mathcal{L}) \left(\begin{array}{l} \widetilde{\omega}_{\mathcal{C}}([l_1, l_2]) \geq \widetilde{\min}\{\widetilde{\omega}_{\mathcal{C}}(l_1), \widetilde{\omega}_{\mathcal{C}}(l_2)\}, \\ \pi_{\mathcal{C}}([l_1, l_2]) \leq \max\{\pi_{\mathcal{C}}(l_1), \pi_{\mathcal{C}}(l_2)\} \end{array} \right).$

Note that for all $l \in L$, $\widetilde{\omega}_{\mathcal{C}}(0) \geq \widetilde{\omega}_{\mathcal{C}}(l)$ and $\pi_{\mathcal{C}}(0) \leq \pi_{\mathcal{C}}(l)$. If $\alpha \neq 0$, $\widetilde{\omega}_{\mathcal{C}}(\alpha l) = \widetilde{\omega}_{\mathcal{C}}(l)$, since $\widetilde{\omega}_{\mathcal{C}}(\alpha l) \geq \widetilde{\omega}_{\mathcal{C}}(l) = \widetilde{\omega}_{\mathcal{C}}(\alpha^{-1}(\alpha l)) \geq \widetilde{\omega}_{\mathcal{C}}(\alpha l)$. Similarly, $\pi_{\mathcal{C}}(\alpha l) = \pi_{\mathcal{C}}(l)$.

Definition 3.2. A CC Lie subalgebra $\mathcal{C} = \langle \widetilde{\omega}_{\mathcal{C}}, \pi_{\mathcal{C}} \rangle$ of \mathcal{L} is a CC Lie ideal of \mathcal{L} if it meets the conditions (A) and (B) of Definition 3.1, respectively, and the condition (E), where

$$(E) (\forall l_1, l_2 \in \mathcal{L}) \left(\begin{array}{l} \widetilde{\omega}_{\mathcal{C}}([l_1, l_2]) \geq \widetilde{\omega}_{\mathcal{C}}(l_1), \\ \pi_{\mathcal{C}}([l_1, l_2]) \leq \pi_{\mathcal{C}}(l_1) \end{array} \right).$$

The Next two examples illustrate the above two definitions.

Example 3.1. Let $\mathcal{L} = \mathbb{R}^3 = \{(n, y, m) : n, y, m \in \mathbb{R}\}$. Then, $(\mathbb{R}^3, [\cdot, \cdot])$ is a real Lie algebra, where the bracket $[\cdot, \cdot]$ is defined as $[l_1, l_2] = l_1 \times l_2$ (the usual cross product). We define a CCS as:

$$\mathcal{C}(n, y, m) = \begin{cases} \langle [0.7, 0.9], -0.8 \rangle, & \text{if } n = y = m; \\ \langle [0.6, 0.8], -0.7 \rangle, & \text{otherwise.} \end{cases}$$

It is clear that, $\mathcal{C} = \langle \widetilde{\omega}_{\mathcal{C}}, \pi_{\mathcal{C}} \rangle$ is a CC Lie subalgebra of \mathbb{R}^3 . But it is not a CC Lie ideal, since $\widetilde{\omega}_{\mathcal{C}}([(1, 1, 1), (1, 2, 4)]) = \widetilde{\omega}_{\mathcal{C}}((2, -3, 1)) = [0.6, 0.8] \not\geq [0.7, 0.9] = \widetilde{\omega}_{\mathcal{C}}((1, 1, 1))$.

Example 3.2. Let $\mathcal{L} = \mathbb{R}^3 = \{(n, y, m) : n, y, m \in \mathbb{R}\}$. Then, $(\mathbb{R}^3, [\cdot, \cdot])$ is a real Lie algebra, where the bracket $[\cdot, \cdot]$ is defined as $[l_1, l_2] = l_1 \times l_2$ (the usual cross product). We define a CCS as:

$$\mathcal{C}(n, y, m) = \begin{cases} \langle [0.7, 0.9], -0.8 \rangle, & \text{if } n = y = m = 0; \\ \langle [0.6, 0.8], -0.7 \rangle, & \text{otherwise.} \end{cases}$$

Clearly, one can obtain that $\mathcal{C} = \langle \widetilde{\omega}_{\mathcal{C}}, \pi_{\mathcal{C}} \rangle$ is a CC Lie ideal of \mathbb{R}^3 .

Theorem 3.3. Any CC Lie ideal is a CC Lie subalgebra.

Proof. The proof is obvious. □

The converse of Theorem 3.3 does not hold in general.

Example 3.3. Let $\mathcal{L} = \mathbb{R}[n]$ be the set of all polynomials with coefficients in \mathbb{R} . Then, $(\mathbb{R}[n], [\cdot, \cdot])$ is a real Lie algebra over \mathbb{R} , where the bracket $[\cdot, \cdot]$ is defined as

$$[p(n), q(n)] = q(0)p(n) - p(0)q(n).$$

We define a CCS as :

$$\mathcal{C}(p(n)) = \left\langle \left[0, \frac{1}{\deg(p(n)) + 1}\right], \frac{-1}{\deg(p(n)) + 2} \right\rangle.$$

Note that, for any $p(n), q(n) \in \mathbb{R}[n]$, and $\alpha \in \mathbb{R}$:

$$\deg(p(n) + q(n)) \leq \max\{\deg(p(n)), \deg(q(n))\},$$

$$\deg(\alpha p(n)) \leq \deg(p(n)),$$

$$\deg([p(n), q(n)]) \leq \max\{\deg(p(n)), \deg(q(n))\}.$$

It is clear that, $\mathcal{C} = \langle \widetilde{\omega}_{\mathcal{C}}, \pi_{\mathcal{C}} \rangle$ is a CC Lie subalgebra of $\mathbb{R}[n]$, but it is not a CC Lie ideal, since

$$\pi_{\mathcal{C}}([n^3 + 1, n^5 + 1]) = \pi_{\mathcal{C}}(n^3 - n^5) = \frac{-1}{7} \not\leq \frac{-1}{5} = \pi_{\mathcal{C}}(n^3 + 1).$$

Theorem 3.4. A CCS $\mathcal{C} = \langle \widetilde{\omega}_{\mathcal{C}}, \pi_{\mathcal{C}} \rangle$ is a CC Lie subalgebra (ideal) if and only if

$$\Upsilon(\widetilde{\omega}_{\mathcal{C}}; \widetilde{\nu}) = \{l \in \mathcal{L} \mid \widetilde{\omega}_{\mathcal{C}}(l) \geq \widetilde{\nu}\} \text{ and } \Upsilon(\pi_{\mathcal{C}}; \gamma) = \{l \in \mathcal{L} \mid \pi_{\mathcal{C}}(l) \leq \gamma\}$$

are Lie subalgebras (ideals) of \mathcal{L} for any $(\widetilde{\nu}, \gamma) \in \text{Im}(\widetilde{\omega}_{\mathcal{C}}) \times \text{Im}(\pi_{\mathcal{C}})$, where $\widetilde{\nu} \in \mathbb{P}[0, 1]$ and $\gamma \in [-1, 0]$.

Proof. Assume that $\mathcal{C} = \langle \widetilde{\omega}_{\mathcal{C}}, \pi_{\mathcal{C}} \rangle$ is a CC Lie subalgebra, and let $\widetilde{\nu} \in \text{Im}(\widetilde{\omega}_{\mathcal{C}})$ and $\gamma \in \text{Im}(\pi_{\mathcal{C}})$. Let $l_1, l_2 \in \Upsilon(\widetilde{\omega}_{\mathcal{C}}; \widetilde{\nu})$ and $\alpha \in F$. Then $\widetilde{\omega}_{\mathcal{C}}(l_1) \geq \widetilde{\nu}$ and $\widetilde{\omega}_{\mathcal{C}}(l_2) \geq \widetilde{\nu}$. It follows that

$$\begin{aligned} \widetilde{\omega}_{\mathcal{C}}(l_1 + l_2) &\geq \widetilde{\min}\{\widetilde{\omega}_{\mathcal{C}}(l_1), \widetilde{\omega}_{\mathcal{C}}(l_2)\} \geq \widetilde{\nu}, \\ \widetilde{\omega}_{\mathcal{C}}(\alpha l_1) &\geq \widetilde{\omega}_{\mathcal{C}}(l_1) \geq \widetilde{\nu}, \\ \widetilde{\omega}_{\mathcal{C}}([l_1, l_2]) &\geq \widetilde{\min}\{\widetilde{\omega}_{\mathcal{C}}(l_1), \widetilde{\omega}_{\mathcal{C}}(l_2)\} \geq \widetilde{\nu} \end{aligned}$$

and hence, $l_1 + l_2, \alpha l_1, [l_1, l_2] \in \Upsilon(\widetilde{\omega}_{\mathcal{C}}; \widetilde{\nu})$. Thus, $\Upsilon(\widetilde{\omega}_{\mathcal{C}}; \widetilde{\nu})$ is a Lie subalgebra of \mathcal{L} . The proof of $\Upsilon(\pi_{\mathcal{C}}; \gamma)$ is similar.

Conversely, suppose that $\Upsilon(\widetilde{\omega}_{\mathcal{C}}; \widetilde{\nu})$ and $\Upsilon(\pi_{\mathcal{C}}; \gamma)$ are Lie subalgebras of \mathcal{L} for any $\widetilde{\nu} \in \text{Im}(\widetilde{\omega}_{\mathcal{C}})$, $\gamma \in \text{Im}(\pi_{\mathcal{C}})$. Let $l_1, l_2 \in \mathcal{L}$ and $\alpha \in F$. Taking

$$\widetilde{\nu} = \widetilde{\min}\{\widetilde{\omega}_{\mathcal{C}}(l_1), \widetilde{\omega}_{\mathcal{C}}(l_2)\}.$$

Thus, $l_1, l_2 \in \Upsilon(\widetilde{\omega}_{\mathcal{C}}; \widetilde{\nu})$. Then, we have $l_1 + l_2 \in \Upsilon(\widetilde{\omega}_{\mathcal{C}}; \widetilde{\nu})$, which means

$$\widetilde{\omega}_{\mathcal{C}}(l_1 + l_2) \geq \widetilde{\nu} = \widetilde{\min}\{\widetilde{\omega}_{\mathcal{C}}(l_1), \widetilde{\omega}_{\mathcal{C}}(l_2)\}.$$

Similarly, we can show that:

$$\begin{aligned} \widetilde{\omega}_{\mathcal{C}}(\alpha l_1) &\geq \widetilde{\omega}_{\mathcal{C}}(l_1), \\ \widetilde{\omega}_{\mathcal{C}}([l_1, l_2]) &\geq \widetilde{\min}\{\widetilde{\omega}_{\mathcal{C}}(l_1), \widetilde{\omega}_{\mathcal{C}}(l_2)\}. \end{aligned}$$

In a similar way, we use the fact that $\Upsilon(\pi_{\mathcal{C}}; \gamma)$ is a Lie subalgebras of \mathcal{L} to show:

$$\begin{aligned} \pi_{\mathcal{C}}(l_1 + l_2) &\leq \max\{\pi_{\mathcal{C}}(l_1), \pi_{\mathcal{C}}(l_2)\}, \\ \pi_{\mathcal{C}}(\alpha l_1) &\leq \pi_{\mathcal{C}}(l_1), \\ \pi_{\mathcal{C}}([l_1, l_2]) &\leq \max\{\pi_{\mathcal{C}}(l_1), \pi_{\mathcal{C}}(l_2)\}. \end{aligned}$$

Hence, $\mathcal{C} = \langle \widetilde{\omega}_{\mathcal{C}}, \pi_{\mathcal{C}} \rangle$ is a CC Lie subalgebra. The proof of the Lie ideal is similar. \square

Theorem 3.5. Let $\mathcal{C} = \langle \widetilde{\omega}_{\mathcal{C}}, \pi_{\mathcal{C}} \rangle$ be a CC Lie subalgebra. Define a binary relation \bowtie on \mathcal{L} by

$$(\forall l_1, l_2 \in \mathcal{L})(l_1 \bowtie l_2) \Leftrightarrow \begin{pmatrix} \widetilde{\omega}_{\mathcal{C}}(l_1 - l_2) = \widetilde{\omega}_{\mathcal{C}}(0), \\ \pi_{\mathcal{C}}(l_1 - l_2) = \pi_{\mathcal{C}}(0) \end{pmatrix}.$$

Then, \bowtie is an equivalence relation on \mathcal{L} .

Proof. Clearly, the reflexivity and symmetry of \bowtie are trivial. So, we only need to demonstrate that \bowtie is transitive. Let $l_1, l_2, l_3 \in \mathcal{L}$. If $l_1 \bowtie l_2$ and $l_2 \bowtie l_3$, then $\widetilde{\omega}_{\mathcal{C}}(l_1 - l_2) = \widetilde{\omega}_{\mathcal{C}}(0)$, $\widetilde{\omega}_{\mathcal{C}}(l_2 - l_3) = \widetilde{\omega}_{\mathcal{C}}(0)$ and $\pi_{\mathcal{C}}(l_1 - l_2) = \pi_{\mathcal{C}}(0)$, $\pi_{\mathcal{C}}(l_2 - l_3) = \pi_{\mathcal{C}}(0)$. Thus, it follows that

$$\widetilde{\omega}_{\mathcal{C}}(0) \geq \widetilde{\omega}_{\mathcal{C}}(l_1 - l_3) = \widetilde{\omega}_{\mathcal{C}}(l_1 - l_2 + l_2 - l_3) \geq \widetilde{\min}\{\widetilde{\omega}_{\mathcal{C}}(l_1 - l_2), \widetilde{\omega}_{\mathcal{C}}(l_2 - l_3)\} = \widetilde{\omega}_{\mathcal{C}}(0)$$

and

$$\pi_{\mathcal{C}}(0) \leq \pi_{\mathcal{C}}(l_1 - l_3) = \pi_{\mathcal{C}}(l_1 - l_2 + l_2 - l_3) \leq \max\{\pi_{\mathcal{C}}(l_1 - l_2), \pi_{\mathcal{C}}(l_2 - l_3)\} = \pi_{\mathcal{C}}(0).$$

Hence, \bowtie is an equivalence relation on \mathcal{L} . \square

Theorem 3.6. Let $\mathcal{H}_0 \subset \mathcal{H}_1 \subset \dots \subset \mathcal{H}_n = \mathcal{L}$ be a chain of Lie algebra \mathcal{L} . Then, there exists a CC Lie subalgebras $\mathcal{C} = \langle \widetilde{\omega}_{\mathcal{C}}, \pi_{\mathcal{C}} \rangle$ for which the level subsets

$$\Upsilon(\widetilde{\omega}_{\mathcal{C}}; \widetilde{\nu}_k) = \{l \in \mathcal{L} \mid \widetilde{\omega}_{\mathcal{C}}(l) \geq \widetilde{\nu}_k\} \text{ and } \Upsilon(\pi_{\mathcal{C}}; \gamma_k) = \{l \in \mathcal{L} \mid \pi_{\mathcal{C}}(l) \leq \gamma_k\},$$

$(\widetilde{\nu}_k, \gamma_k) \in \text{Im}(\widetilde{\omega}_{\mathcal{C}}) \times \text{Im}(\pi_{\mathcal{C}})$, where $\widetilde{\nu}_k \in \mathbb{P}[0, 1]$ and $\gamma_k \in [-1, 0]$, $1 \leq k \leq n$ coincide with this chain.

Proof. Take $\widetilde{\nu}_k$ and γ_k , $k = 0, 1, \dots, n$ such that

$$\widetilde{\nu}_0 \geq \widetilde{\nu}_1 \geq \dots \geq \widetilde{\nu}_n$$

and

$$\gamma_0 \leq \gamma_1 \leq \dots \leq \gamma_n.$$

We can choose $\widetilde{\nu}_k = [0, \frac{1}{1+k}]$ and $\gamma_k = \frac{-1}{1+k}$, $k = 0, 1, \dots, n$. Let $\mathcal{C} = \langle \widetilde{\omega}_{\mathcal{C}}, \pi_{\mathcal{C}} \rangle$ be a CCS of \mathcal{L} defined by

$$\mathcal{C}(l) = \begin{cases} \langle \widetilde{\nu}_0, \gamma_0 \rangle, & \text{if } l \in \mathcal{H}_0; \\ \langle \widetilde{\nu}_1, \gamma_1 \rangle, & \text{if } l \in \mathcal{H}_1 - \mathcal{H}_0; \\ \langle \widetilde{\nu}_2, \gamma_2 \rangle, & \text{if } l \in \mathcal{H}_2 - \mathcal{H}_1; \\ \vdots & \\ \langle \widetilde{\nu}_n, \gamma_n \rangle, & \text{if } l \in \mathcal{H}_n - \mathcal{H}_{n-1}. \end{cases}$$

We first prove that $\mathcal{C} = \langle \widetilde{\omega}_{\mathcal{C}}, \pi_{\mathcal{C}} \rangle$ is a CC Lie subalgebra of \mathcal{L} . Let $l_1, l_2 \in \mathcal{L}$ and $\alpha \in F$. Suppose that $l_1 \in \mathcal{H}_i - \mathcal{H}_{i-1}$ and $l_2 \in \mathcal{H}_j - \mathcal{H}_{j-1}$ for some i, j . We may assume that $j \leq i$. Then, we have $l_1, l_2 \in \mathcal{H}_i$. Thus, $l_1 + l_2, \alpha l_1, [l_1, l_2] \in \mathcal{H}_i$. It follows that

$$\begin{aligned} \widetilde{\omega}_{\mathcal{C}}(l_1 + l_2) &\geq \widetilde{\nu}_i = \widetilde{\min}\{\widetilde{\nu}_i, \widetilde{\nu}_j\} = \widetilde{\min}\{\widetilde{\omega}_{\mathcal{C}}(l_1), \widetilde{\omega}_{\mathcal{C}}(l_2)\}, \\ \widetilde{\omega}_{\mathcal{C}}(\alpha l_1) &\geq \widetilde{\nu}_i = \widetilde{\omega}_{\mathcal{C}}(l_1), \\ \widetilde{\omega}_{\mathcal{C}}([l_1, l_2]) &\geq \widetilde{\nu}_i = \widetilde{\min}\{\widetilde{\nu}_i, \widetilde{\nu}_j\} = \widetilde{\min}\{\widetilde{\omega}_{\mathcal{C}}(l_1), \widetilde{\omega}_{\mathcal{C}}(l_2)\}, \end{aligned}$$

and

$$\begin{aligned} \pi_{\mathcal{C}}(l_1 + l_2) &\leq \gamma_i = \max\{\gamma_i, \gamma_j\} = \max\{\pi_{\mathcal{C}}(l_1), \pi_{\mathcal{C}}(l_2)\}, \\ \pi_{\mathcal{C}}(\alpha l_1) &\leq \gamma_i = \pi_{\mathcal{C}}(l_1), \\ \pi_{\mathcal{C}}([l_1, l_2]) &\leq \gamma_i = \max\{\gamma_i, \gamma_j\} = \max\{\pi_{\mathcal{C}}(l_1), \pi_{\mathcal{C}}(l_2)\}. \end{aligned}$$

So, $\mathcal{C} = \langle \widetilde{\omega}_{\mathcal{C}}, \pi_{\mathcal{C}} \rangle$ is a CC Lie subalgebra of \mathcal{L} and all its level sets are Lie algebras, since $\text{Im}(\widetilde{\omega}_{\mathcal{C}}) = \{\widetilde{\nu}_0, \widetilde{\nu}_1, \dots, \widetilde{\nu}_n\}$ and $\text{Im}(\pi_{\mathcal{C}}) = \{\gamma_0, \gamma_1, \dots, \gamma_n\}$ are level subsets of \mathcal{C} form chains:

$$\Upsilon(\widetilde{\omega}_{\mathcal{C}}; \widetilde{\nu}_0) \subset \Upsilon(\widetilde{\omega}_{\mathcal{C}}; \widetilde{\nu}_1) \subset \dots \subset \Upsilon(\widetilde{\omega}_{\mathcal{C}}; \widetilde{\nu}_n) = \mathcal{L},$$

and

$$\Upsilon(\pi_{\mathcal{C}}; \gamma_0) \subset \Upsilon(\pi_{\mathcal{C}}; \gamma_1) \subset \dots \subset \Upsilon(\pi_{\mathcal{C}}; \gamma_n) = \mathcal{L},$$

respectively. We now prove that

$$\Upsilon(\widetilde{\omega}_{\mathcal{C}}; \widetilde{\nu}_k) = \mathcal{H}_k = \Upsilon(\pi_{\mathcal{C}}; \gamma_k), k = 0, 1, \dots, n.$$

Clearly, $\mathcal{H}_k \subseteq \Upsilon(\widetilde{\omega}_{\mathcal{C}}; \widetilde{\nu}_k)$ and $\mathcal{H}_k \subseteq \Upsilon(\pi_{\mathcal{C}}; \gamma_k)$. If $l \in \Upsilon(\widetilde{\omega}_{\mathcal{C}}; \widetilde{\nu}_k)$, then $\widetilde{\omega}_{\mathcal{C}}(l) \geq \widetilde{\nu}_k$ and so $\widetilde{\omega}_{\mathcal{C}}(l) = \widetilde{\nu}_i$ for some $i \leq k$. Thus, $l \in \mathcal{H}_i$. Since $\mathcal{H}_i \subseteq \mathcal{H}_k$, it follows that $l \in \mathcal{H}_k$. Consequently, $\mathcal{H}_k = \Upsilon(\widetilde{\omega}_{\mathcal{C}}; \widetilde{\nu}_k)$. The proof of $\Upsilon(\pi_{\mathcal{C}}; \gamma)$ is similar. \square

Definition 3.7. Let $\mathcal{C} = \langle \widetilde{\omega}_{\mathcal{C}}, \pi_{\mathcal{C}} \rangle$ and $\mathcal{D} = \langle \widetilde{\omega}_{\mathcal{D}}, \pi_{\mathcal{D}} \rangle$ be two CCSs on \mathcal{L} . If $\mathcal{C} = \langle \widetilde{\omega}_{\mathcal{C}}, \pi_{\mathcal{C}} \rangle$ is a CC relation on \mathcal{L} , then $\mathcal{C} = \langle \widetilde{\omega}_{\mathcal{C}}, \pi_{\mathcal{C}} \rangle$ is called a CC relation on $\mathcal{D} = \langle \widetilde{\omega}_{\mathcal{D}}, \pi_{\mathcal{D}} \rangle$ if $\widetilde{\omega}_{\mathcal{C}}(l_1, l_2) \leq \widetilde{\min}\{\widetilde{\omega}_{\mathcal{D}}(l_1), \widetilde{\omega}_{\mathcal{D}}(l_2)\}$ and $\widetilde{\pi}_{\mathcal{C}}(l_1, l_2) \geq \widetilde{\max}\{\widetilde{\pi}_{\mathcal{D}}(l_1), \widetilde{\pi}_{\mathcal{D}}(l_2)\}$ for all $l_1, l_2 \in \mathcal{L}$.

Definition 3.8. Let $\mathcal{C} = \langle \widetilde{\omega}_{\mathcal{C}}, \pi_{\mathcal{C}} \rangle$ and $\mathcal{D} = \langle \widetilde{\omega}_{\mathcal{D}}, \pi_{\mathcal{D}} \rangle$ be two CCSs on \mathcal{L} . Then, the generalized Cartesian product $\mathcal{C} \times \mathcal{D}$ is presented as:

$$\begin{aligned} \mathcal{C} \times \mathcal{D} &= \langle \widetilde{\omega}_{\mathcal{C}}, \pi_{\mathcal{C}} \rangle \times \langle \widetilde{\omega}_{\mathcal{D}}, \pi_{\mathcal{D}} \rangle \\ &= \langle \widetilde{\omega}_{\mathcal{C}} \times \widetilde{\omega}_{\mathcal{D}}, \pi_{\mathcal{C}} \times \pi_{\mathcal{D}} \rangle, \end{aligned}$$

where $(\widetilde{\omega}_{\mathcal{C}} \times \widetilde{\omega}_{\mathcal{D}})(l_1, l_2) = \widetilde{\min}\{\widetilde{\omega}_{\mathcal{C}}(l_1), \widetilde{\omega}_{\mathcal{D}}(l_2)\}$ and $(\pi_{\mathcal{C}} \times \pi_{\mathcal{D}})(l_1, l_2) = \widetilde{\max}\{\widetilde{\pi}_{\mathcal{C}}(l_1), \widetilde{\pi}_{\mathcal{D}}(l_2)\}$ for all $(l_1, l_2) \in \mathcal{L} \times \mathcal{L}$.

It is clear that the generalized Cartesian product $\mathcal{C} \times \mathcal{D}$ is always a CC structure on $\mathcal{L} \times \mathcal{L}$.

The proof of the next theorem is trivial, so it is omitted.

Theorem 3.9. Let $\mathcal{C} = \langle \widetilde{\omega}_{\mathcal{C}}, \pi_{\mathcal{C}} \rangle$ and $\mathcal{D} = \langle \widetilde{\omega}_{\mathcal{D}}, \pi_{\mathcal{D}} \rangle$ be two CCSs on \mathcal{L} . Then,

(1) $\mathcal{C} \times \mathcal{D}$ is a CC relation on \mathcal{L} .

(2) $\Upsilon(\widetilde{\omega}_{\mathcal{C}} \times \widetilde{\omega}_{\mathcal{D}}; \widetilde{\nu}) = \Upsilon(\widetilde{\omega}_{\mathcal{C}}; \widetilde{\nu}) \times \Upsilon(\widetilde{\omega}_{\mathcal{D}}; \widetilde{\nu})$ and $\Upsilon(\pi_{\mathcal{C}} \times \pi_{\mathcal{D}}; \gamma) = \Upsilon(\pi_{\mathcal{C}}; \gamma) \times \Upsilon(\pi_{\mathcal{D}}; \gamma)$

$\forall \widetilde{\nu} \in \mathbb{P}[0, 1], \gamma \in [-1, 0]$.

Theorem 3.10. Let $\mathcal{C} = \langle \widetilde{\omega}_{\mathcal{C}}, \pi_{\mathcal{C}} \rangle$ and $\mathcal{D} = \langle \widetilde{\omega}_{\mathcal{D}}, \pi_{\mathcal{D}} \rangle$ be two CC Lie subalgebras on \mathcal{L} . Then, $\mathcal{C} \times \mathcal{D}$ is a CC Lie subalgebra on $\mathcal{L} \times \mathcal{L}$.

Proof. Let $l = (l_1, l_2), n = (n_1, n_2) \in \mathcal{L} \times \mathcal{L}$. Then,

$$\begin{aligned} (\widetilde{\omega}_{\mathcal{C}} \times \widetilde{\omega}_{\mathcal{D}})(l + n) &= (\widetilde{\omega}_{\mathcal{C}} \times \widetilde{\omega}_{\mathcal{D}})((l_1, l_2) + (n_1, n_2)) \\ &= (\widetilde{\omega}_{\mathcal{C}} \times \widetilde{\omega}_{\mathcal{D}})((l_1 + n_1), (l_2 + n_2)) \\ &= \widetilde{\min}\{\widetilde{\omega}_{\mathcal{C}}(l_1 + n_1), \widetilde{\omega}_{\mathcal{D}}(l_2 + n_2)\} \\ &\geq \widetilde{\min}\{\widetilde{\min}\{\widetilde{\omega}_{\mathcal{C}}(l_1), \widetilde{\omega}_{\mathcal{C}}(n_1)\}, \widetilde{\min}\{\widetilde{\omega}_{\mathcal{D}}(l_2), \widetilde{\omega}_{\mathcal{D}}(n_2)\}\} \\ &= \widetilde{\min}\{\widetilde{\min}\{\widetilde{\omega}_{\mathcal{C}}(l_1), \widetilde{\omega}_{\mathcal{D}}(l_2)\}, \widetilde{\min}\{\widetilde{\omega}_{\mathcal{C}}(n_1), \widetilde{\omega}_{\mathcal{D}}(n_2)\}\} \\ &= \widetilde{\min}\{(\widetilde{\omega}_{\mathcal{C}} \times \widetilde{\omega}_{\mathcal{D}})(l_1, l_2), (\widetilde{\omega}_{\mathcal{C}} \times \widetilde{\omega}_{\mathcal{D}})(n_1, n_2)\} \\ &= \widetilde{\min}\{(\widetilde{\omega}_{\mathcal{C}} \times \widetilde{\omega}_{\mathcal{D}})(l), (\widetilde{\omega}_{\mathcal{C}} \times \widetilde{\omega}_{\mathcal{D}})(n)\}, \end{aligned}$$

$$\begin{aligned}
(\widetilde{\omega}_{\mathcal{C}} \times \widetilde{\omega}_{\mathcal{D}})(\alpha l) &= (\widetilde{\omega}_{\mathcal{C}} \times \widetilde{\omega}_{\mathcal{D}})(\alpha(l_1, l_2)) \\
&= (\widetilde{\omega}_{\mathcal{C}} \times \widetilde{\omega}_{\mathcal{D}})(\alpha l_1, \alpha l_2) \\
&= \widetilde{\min}\{\widetilde{\omega}_{\mathcal{C}}(\alpha l_1), \widetilde{\omega}_{\mathcal{D}}(\alpha l_2)\} \\
&\geq \widetilde{\min}\{\widetilde{\omega}_{\mathcal{C}}(l_1), \widetilde{\omega}_{\mathcal{D}}(l_2)\} \\
&= (\widetilde{\omega}_{\mathcal{C}} \times \widetilde{\omega}_{\mathcal{D}})(l_1, l_2) \\
&= (\widetilde{\omega}_{\mathcal{C}} \times \widetilde{\omega}_{\mathcal{D}})(l),
\end{aligned}$$

$$\begin{aligned}
(\widetilde{\omega}_{\mathcal{C}} \times \widetilde{\omega}_{\mathcal{D}})([l + n]) &= (\widetilde{\omega}_{\mathcal{C}} \times \widetilde{\omega}_{\mathcal{D}})([(l_1, l_2) + (n_1, n_2)]) \\
&\geq \widetilde{\min}\{\widetilde{\min}\{\widetilde{\omega}_{\mathcal{C}}(l_1), \widetilde{\omega}_{\mathcal{D}}(l_2)\}, \widetilde{\min}\{\widetilde{\omega}_{\mathcal{C}}(n_1), \widetilde{\omega}_{\mathcal{D}}(n_2)\}\} \\
&= \widetilde{\min}\{(\widetilde{\omega}_{\mathcal{C}} \times \widetilde{\omega}_{\mathcal{D}})(l_1, l_2), (\widetilde{\omega}_{\mathcal{C}} \times \widetilde{\omega}_{\mathcal{D}})(n_1, n_2)\} \\
&= \widetilde{\min}\{(\widetilde{\omega}_{\mathcal{C}} \times \widetilde{\omega}_{\mathcal{D}})(l), (\widetilde{\omega}_{\mathcal{C}} \times \widetilde{\omega}_{\mathcal{D}})(n)\}.
\end{aligned}$$

Also,

$$\begin{aligned}
(\pi_{\mathcal{C}} \times \pi_{\mathcal{D}})(l + n) &= (\pi_{\mathcal{C}} \times \pi_{\mathcal{D}})((l_1, l_2) + (n_1, n_2)) \\
&= (\pi_{\mathcal{C}} \times \pi_{\mathcal{D}})((l_1 + n_1), (l_2 + n_2)) \\
&= \max\{\pi_{\mathcal{C}}(l_1 + n_1), \pi_{\mathcal{D}}(l_2 + n_2)\} \\
&\leq \max\{\max\{\pi_{\mathcal{C}}(l_1), \pi_{\mathcal{C}}(n_1)\}, \max\{\pi_{\mathcal{D}}(l_2), \pi_{\mathcal{D}}(n_2)\}\} \\
&= \max\{\max\{\pi_{\mathcal{C}}(l_1), \pi_{\mathcal{D}}(l_2)\}, \max\{\pi_{\mathcal{C}}(n_1), \pi_{\mathcal{D}}(n_2)\}\} \\
&= \max\{(\pi_{\mathcal{C}} \times \pi_{\mathcal{D}})(l_1, l_2), (\pi_{\mathcal{C}} \times \pi_{\mathcal{D}})(n_1, n_2)\} \\
&= \max\{(\pi_{\mathcal{C}} \times \pi_{\mathcal{D}})(l), (\pi_{\mathcal{C}} \times \pi_{\mathcal{D}})(n)\},
\end{aligned}$$

$$\begin{aligned}
(\pi_{\mathcal{C}} \times \pi_{\mathcal{D}})(\alpha l) &= (\pi_{\mathcal{C}} \times \pi_{\mathcal{D}})(\alpha(l_1, l_2)) \\
&= (\pi_{\mathcal{C}} \times \pi_{\mathcal{D}})(\alpha l_1, \alpha l_2) \\
&= \max\{\pi_{\mathcal{C}}(\alpha l_1), \pi_{\mathcal{D}}(\alpha l_2)\} \\
&\leq \max\{\pi_{\mathcal{C}}(l_1), \pi_{\mathcal{D}}(l_2)\} \\
&= (\pi_{\mathcal{C}} \times \pi_{\mathcal{D}})(l_1, l_2) \\
&= (\pi_{\mathcal{C}} \times \pi_{\mathcal{D}})(l)
\end{aligned}$$

and

$$\begin{aligned}
(\pi_{\mathcal{C}} \times \pi_{\mathcal{D}})([l + n]) &= (\pi_{\mathcal{C}} \times \pi_{\mathcal{D}})([(l_1, l_2) + (n_1, n_2)]) \\
&\leq \max\{\max\{\pi_{\mathcal{C}}(l_1), \pi_{\mathcal{D}}(l_2)\}, \max\{\pi_{\mathcal{C}}(n_1), \pi_{\mathcal{D}}(n_2)\}\} \\
&= \max\{(\pi_{\mathcal{C}} \times \pi_{\mathcal{D}})(l_1, l_2), (\pi_{\mathcal{C}} \times \pi_{\mathcal{D}})(n_1, n_2)\} \\
&= \max\{(\pi_{\mathcal{C}} \times \pi_{\mathcal{D}})(l), (\pi_{\mathcal{C}} \times \pi_{\mathcal{D}})(n)\}.
\end{aligned}$$

Hence, $\mathcal{C} \times \mathcal{D}$ is a CC Lie subalgebra on $\mathcal{L} \times \mathcal{L}$. □

Definition 3.11. Let $\mathcal{C} = \langle \widetilde{\omega}_{\mathcal{C}}, \pi_{\mathcal{C}} \rangle$ and $\mathcal{D} = \langle \widetilde{\omega}_{\mathcal{D}}, \pi_{\mathcal{D}} \rangle$ be two CC Lie ideals of \mathcal{L} . Then, \mathcal{C} is the same type of \mathcal{D} if $\mathcal{C} = \mathcal{D} \circ \Psi$ for some $\Psi \in \text{Aut } \mathcal{L}$, i.e., $\widetilde{\omega}_{\mathcal{C}}(l) = \widetilde{\omega}_{\mathcal{D}}(\Psi(l))$ and $\pi_{\mathcal{C}}(l) = \pi_{\mathcal{D}}(\Psi(l))$ for all $l \in \mathcal{L}$.

Theorem 3.12. Let $\mathcal{C} = \langle \widetilde{\omega}_{\mathcal{C}}, \pi_{\mathcal{C}} \rangle$ and $\mathcal{D} = \langle \widetilde{\omega}_{\mathcal{D}}, \pi_{\mathcal{D}} \rangle$ be two CC Lie ideals of \mathcal{L} . Then, \mathcal{C} is isomorphic to \mathcal{D} if and only if \mathcal{C} is a CC Lie ideal having the same type of \mathcal{D} .

Proof. Let $\mathcal{C} = \langle \widetilde{\omega}_{\mathcal{C}}, \pi_{\mathcal{C}} \rangle$ and $\mathcal{D} = \langle \widetilde{\omega}_{\mathcal{D}}, \pi_{\mathcal{D}} \rangle$ be two CC Lie ideals of \mathcal{L} . If \mathcal{C} is isomorphic to \mathcal{D} , then clearly \mathcal{C} is a crossing Lie ideal having the same type of \mathcal{D} . Now, suppose that \mathcal{C} is a crossing Lie ideal having the same type of \mathcal{D} . Then, $\exists \Psi \in \text{Aut}(\mathcal{L})$ such that

$$\widetilde{\omega}_{\mathcal{C}}(l_1) = \widetilde{\omega}_{\mathcal{C}}(\Psi(l_1)) \text{ and } \pi_{\mathcal{C}}(l_1) = \pi_{\mathcal{C}}(\Psi(l_1))$$

for any $l_1 \in \mathcal{L}$. Let $\Omega : \mathcal{C}(\mathcal{L}) \rightarrow \mathcal{D}(\mathcal{L})$ be a map presented by:

$$\Omega(\mathcal{C}(l_1)) = \mathcal{D}(\Psi(l_1))$$

for any $l_1 \in \mathcal{L}$. That is,

$$\Omega(\widetilde{\omega}_{\mathcal{C}}(l_1)) = \widetilde{\omega}_{\mathcal{D}}(\Psi(l_1)) \text{ and } \Omega(\pi_{\mathcal{C}}(l_1)) = \pi_{\mathcal{D}}(\Psi(l_1))$$

for any $l_1 \in \mathcal{L}$. Then, it is obvious that Ω is surjective. Since $\Omega(\widetilde{\omega}_{\mathcal{C}}(l_1)) = \Omega(\widetilde{\omega}_{\mathcal{C}}(l_2))$ for any $l_1, l_2 \in \mathcal{L}$, then $\widetilde{\omega}_{\mathcal{D}}(\Psi(l_1)) = \widetilde{\omega}_{\mathcal{D}}(\Psi(l_2))$ and therefore $\widetilde{\omega}_{\mathcal{C}}(l_1) = \widetilde{\omega}_{\mathcal{C}}(l_2)$. Similarly, $\Omega(\pi_{\mathcal{C}}(l_1)) = \Omega(\pi_{\mathcal{C}}(\Psi(l_2)))$ for any $l_1, l_2 \in \mathcal{L}$, then $\widetilde{\omega}_{\mathcal{D}}(\Psi(l_1)) = \pi_{\mathcal{D}}(\Psi(l_2))$ and therefore $\pi_{\mathcal{C}}(l_1) = \pi_{\mathcal{C}}(l_2)$. Now, for any $l_1, l_2 \in \mathcal{L}$, we have

$$\Omega(\widetilde{\omega}_{\mathcal{C}}(l_1 + l_2)) = \widetilde{\omega}_{\mathcal{D}}(\Psi(l_1 + l_2)) = \widetilde{\omega}_{\mathcal{D}}(\Psi(l_1) + \Psi(l_2)),$$

$$\Omega(\widetilde{\omega}_{\mathcal{C}}(\alpha l_1)) = \widetilde{\omega}_{\mathcal{D}}(\Psi(\alpha l_1)) = \alpha \widetilde{\omega}_{\mathcal{D}}(\Psi(l_1))$$

and

$$\Omega(\widetilde{\omega}_{\mathcal{C}}([l_1 + l_2])) = \widetilde{\omega}_{\mathcal{D}}(\Psi([l_1 + l_2])) = \widetilde{\omega}_{\mathcal{D}}([\Psi(l_1) + \Psi(l_2)]).$$

Similarly,

$$\Omega(\pi_{\mathcal{C}}(l_1 + l_2)) = \pi_{\mathcal{D}}(\Psi(l_1 + l_2)) = \pi_{\mathcal{D}}(\Psi(l_1) + \Psi(l_2)),$$

$$\pi_{\mathcal{C}}(\alpha l_1) = \pi_{\mathcal{D}}(\Psi(\alpha l_1)) = \alpha \pi_{\mathcal{D}}(\Psi(l_1)),$$

and

$$\Omega(\pi_{\mathcal{C}}([l_1 + l_2])) = \pi_{\mathcal{D}}(\Psi([l_1 + l_2])) = \pi_{\mathcal{D}}([\Psi(l_1) + \Psi(l_2)]).$$

Therefore, Ω is a homomorphism. Thus, \mathcal{C} is isomorphic to \mathcal{D} . □

Theorem 3.13. Let $\mathcal{C} = \langle \widetilde{\omega}_{\mathcal{C}}, \pi_{\mathcal{C}} \rangle$ and $\mathcal{D} = \langle \widetilde{\omega}_{\mathcal{D}}, \pi_{\mathcal{D}} \rangle$ be a CC Lie ideals of \mathcal{L} . If \mathcal{C} is a CC Lie ideal having the same type of \mathcal{D} , then $\Psi \in \text{Aut}(\mathcal{L})$ such that $\widetilde{\omega}_{\mathcal{C}} \circ \Psi = \widetilde{\omega}_{\mathcal{D}}$ and $\pi_{\mathcal{C}} \circ \Psi = \pi_{\mathcal{D}}$.

Proof. The proof is obvious from the Definition 3.11. □

Definition 3.14. Let $\Psi : \mathcal{L}_1 \rightarrow \mathcal{L}_2$ be a homomorphism Lie algebras. For any CC structure $\mathcal{C} = \langle \widetilde{\omega}_{\mathcal{C}}, \pi_{\mathcal{C}} \rangle$ on \mathcal{L}_2 , we present a CC structure $\mathcal{C}^{\Psi} = \langle \widetilde{\omega}_{\mathcal{C}}^{\Psi}, \pi_{\mathcal{C}}^{\Psi} \rangle$ on \mathcal{L}_1 , where $\mathcal{C}^{\Psi}(l) = \mathcal{C}(\Psi(l))$ for any $l \in \mathcal{L}_1$. That is, $\widetilde{\omega}_{\mathcal{C}}^{\Psi}(l) = \widetilde{\omega}_{\mathcal{C}}(\Psi(l))$ and $\pi_{\mathcal{C}}^{\Psi}(l) = \pi_{\mathcal{C}}(\Psi(l))$ for any $l \in \mathcal{L}_1$.

Theorem 3.15. Let $\Psi : \mathcal{L}_1 \rightarrow \mathcal{L}_2$ be a homomorphism Lie algebras. If $\mathcal{C} = \langle \widetilde{\omega}_{\mathcal{C}}, \pi_{\mathcal{C}} \rangle$ is a CC Lie ideal of \mathcal{L}_2 , then $\mathcal{C}^{\Psi} = \langle \widetilde{\omega}_{\mathcal{C}}^{\Psi}, \pi_{\mathcal{C}}^{\Psi} \rangle$ is a CC Lie ideal of \mathcal{L}_1 .

Proof. Let $l_1, l_2 \in \mathcal{L}_1$ and $\alpha \in F$. Then,

$$\begin{aligned} \widetilde{\omega}_{\mathcal{C}}^{\Psi}(l_1 + l_2) &= \widetilde{\omega}_{\mathcal{C}}(\Psi(l_1 + l_2)) = \widetilde{\omega}_{\mathcal{C}}(\Psi(l_1) + \Psi(l_2)) \\ &\geq \widetilde{\min}\{\widetilde{\omega}_{\mathcal{C}}(\Psi(l_1)), \widetilde{\omega}_{\mathcal{C}}(\Psi(l_2))\} \\ &= \widetilde{\min}\{\widetilde{\omega}_{\mathcal{C}}^{\Psi}(l_1), \widetilde{\omega}_{\mathcal{C}}^{\Psi}(l_2)\}, \end{aligned}$$

$$\widetilde{\omega}_{\mathcal{C}}^{\Psi}(\alpha l_1) = \widetilde{\omega}_{\mathcal{C}}(\Psi(\alpha l_1)) = \widetilde{\omega}_{\mathcal{C}}(\alpha \Psi(l_1)) \geq \widetilde{\omega}_{\mathcal{C}}(\Psi(l_1)) = \widetilde{\omega}_{\mathcal{C}}^{\Psi}(l_1)$$

and

$$\widetilde{\omega}_{\mathcal{C}}^{\Psi}([l_1 + l_2]) = \widetilde{\omega}_{\mathcal{C}}(\Psi([l_1 + l_2])) = \widetilde{\omega}_{\mathcal{C}}([\Psi(l_1) + \Psi(l_2)]) \geq \widetilde{\omega}_{\mathcal{C}}(\Psi(l_1)) = \widetilde{\omega}_{\mathcal{C}}^{\Psi}(l_1).$$

Similarly, $\pi_{\mathcal{C}}^{\Psi}(l_1 + l_2) \leq \max\{\pi_{\mathcal{C}}^{\Psi}(l_1), \pi_{\mathcal{C}}^{\Psi}(l_2)\}$, $\pi_{\mathcal{C}}^{\Psi}(\alpha l_1) \leq \pi_{\mathcal{C}}^{\Psi}(l_1)$ and $\widetilde{\omega}_{\mathcal{C}}^{\Psi}([l_1 + l_2]) \leq \widetilde{\omega}_{\mathcal{C}}^{\Psi}(l_1)$. Thus, $\mathcal{C}^{\Psi} = \langle \widetilde{\omega}_{\mathcal{C}}^{\Psi}, \pi_{\mathcal{C}}^{\Psi} \rangle$ is a CC Lie ideal of \mathcal{L}_1 . \square

Theorem 3.16. Let $\Psi : \mathcal{L}_1 \rightarrow \mathcal{L}_2$ be an epimorphism of Lie algebras. If $\mathcal{C}^{\Psi} = \langle \widetilde{\omega}_{\mathcal{C}}^{\Psi}, \pi_{\mathcal{C}}^{\Psi} \rangle$ is a CC Lie ideal of \mathcal{L}_1 , then $\mathcal{C} = \langle \widetilde{\omega}_{\mathcal{C}}, \pi_{\mathcal{C}} \rangle$ is a CC Lie ideal of \mathcal{L}_2 .

Proof. Since Ψ is surjective, then for any $l_1, l_2 \in \mathcal{L}_2$, there are $n_1, n_2 \in \mathcal{L}_1$ such that $l_1 = \Psi(n_1)$ and $l_2 = \Psi(n_2)$. Hence, $\widetilde{\omega}_{\mathcal{C}}(l_1) = \widetilde{\omega}_{\mathcal{C}}^{\Psi}(n_1)$, $\widetilde{\omega}_{\mathcal{C}}(l_2) = \widetilde{\omega}_{\mathcal{C}}^{\Psi}(n_2)$, $\pi_{\mathcal{C}}(l_1) = \pi_{\mathcal{C}}^{\Psi}(n_1)$ and $\pi_{\mathcal{C}}(l_2) = \pi_{\mathcal{C}}^{\Psi}(n_2)$. Now,

$$\begin{aligned} \widetilde{\omega}_{\mathcal{C}}(l_1 + l_2) &= \widetilde{\omega}_{\mathcal{C}}(\Psi(n_1) + \Psi(n_2)) = \widetilde{\omega}_{\mathcal{C}}(\Psi(n_1 + n_2)) \\ &= \widetilde{\omega}_{\mathcal{C}}^{\Psi}(n_1 + n_2) \\ &\geq \widetilde{\min}\{\widetilde{\omega}_{\mathcal{C}}^{\Psi}(n_1), \widetilde{\omega}_{\mathcal{C}}^{\Psi}(n_2)\} \\ &= \widetilde{\min}\{\widetilde{\omega}_{\mathcal{C}}(l_1), \widetilde{\omega}_{\mathcal{C}}(l_2)\}, \end{aligned}$$

$$\begin{aligned} \widetilde{\omega}_{\mathcal{C}}(\alpha l_1) &= \widetilde{\omega}_{\mathcal{C}}(\alpha \Psi(n_1)) = \widetilde{\omega}_{\mathcal{C}}(\Psi(\alpha n_1)) \\ &\geq \widetilde{\omega}_{\mathcal{C}}^{\Psi}(n_1) \\ &= \widetilde{\omega}_{\mathcal{C}}(l_1) \end{aligned}$$

and

$$\begin{aligned} \widetilde{\omega}_{\mathcal{C}}([l_1 + l_2]) &= \widetilde{\omega}_{\mathcal{C}}([\Psi(n_1) + \Psi(n_2)]) = \widetilde{\omega}_{\mathcal{C}}^{\Psi}([n_1 + n_2]) \\ &\geq \widetilde{\omega}_{\mathcal{C}}^{\Psi}(n_1) \\ &= \widetilde{\omega}_{\mathcal{C}}(l_1). \end{aligned}$$

Similarly, $\pi_{\mathcal{C}}(l_1 + l_2) \leq \max\{\pi_{\mathcal{C}}(l_1), \pi_{\mathcal{C}}(l_2)\}$, $\pi_{\mathcal{C}}(\alpha l_1) \leq \pi_{\mathcal{C}}(l_1)$ and $\pi_{\mathcal{C}}([l_1 + l_2]) \leq \pi_{\mathcal{C}}(l_1)$. Thus, $\mathcal{C} = \langle \widetilde{\omega}_{\mathcal{C}}, \pi_{\mathcal{C}} \rangle$ is a CC Lie ideal of \mathcal{L}_2 . \square

In Theorems 3.17–3.20 below, let $\mathcal{C} = \langle \widetilde{\omega}_{\mathcal{C}}, \pi_{\mathcal{C}} \rangle$ and $\mathcal{D} = \langle \widetilde{\omega}_{\mathcal{D}}, \pi_{\mathcal{D}} \rangle$ be two CC Lie ideals of \mathcal{L} .

Theorem 3.17. *If $\widetilde{\omega}_{\mathcal{C}} \circ \Psi = \widetilde{\omega}_{\mathcal{D}}$ and $\pi_{\mathcal{C}} \circ \Psi = \pi_{\mathcal{D}}$ for some $\Psi \in \text{Aut}(\mathcal{L})$, then $g(\widetilde{\omega}_{\mathcal{C}}) = \widetilde{\omega}_{\mathcal{D}}$ and $g(\pi_{\mathcal{C}}) = \pi_{\mathcal{D}}$.*

Proof. Let $\mathcal{C} = \langle \widetilde{\omega}_{\mathcal{C}}, \pi_{\mathcal{C}} \rangle$ and $\mathcal{D} = \langle \widetilde{\omega}_{\mathcal{D}}, \pi_{\mathcal{D}} \rangle$ be CC Lie ideals of \mathcal{L} . Assume that $\exists \Psi \in \text{Aut}(\mathcal{L})$ such that $\widetilde{\omega}_{\mathcal{C}} \circ \Psi = \widetilde{\omega}_{\mathcal{D}}$ and $\pi_{\mathcal{C}} \circ \Psi = \pi_{\mathcal{D}}$, then $\widetilde{\omega}_{\mathcal{C}}(\Psi(l_1)) = \widetilde{\omega}_{\mathcal{D}}(l_1)$ and $\pi_{\mathcal{C}}(\Psi(l_1)) = \pi_{\mathcal{D}}(l_1)$. Thus,

$$\Psi^{-1}(\widetilde{\omega}_{\mathcal{C}}(l_1)) = \sup_{l_2 \in \Psi(l_1)} \widetilde{\omega}_{\mathcal{C}}(l_2) = \widetilde{\omega}_{\mathcal{C}}(\Psi(l_1)) = \widetilde{\omega}_{\mathcal{D}}(l_1)$$

and

$$\Psi^{-1}(\pi_{\mathcal{C}}(l_1)) = \inf_{l_2 \in \Psi(l_1)} \pi_{\mathcal{C}}(l_2) = \pi_{\mathcal{C}}(\Psi(l_1)) = \pi_{\mathcal{D}}(l_1)$$

for all $l_1 \in \mathcal{L}$. If $g = \Psi^{-1}$, then $g \in \text{Aut}(\mathcal{L})$, therefore $g(\widetilde{\omega}_{\mathcal{C}}) = \widetilde{\omega}_{\mathcal{D}}$ and $g(\pi_{\mathcal{C}}) = \pi_{\mathcal{D}}$. \square

Theorem 3.18. *If $g(\widetilde{\omega}_{\mathcal{C}}) = \widetilde{\omega}_{\mathcal{D}}$ and $g(\pi_{\mathcal{C}}) = \pi_{\mathcal{D}}$ for some $g \in \text{Aut}(\mathcal{L})$, then $\exists h \in \text{Aut}(\mathcal{L})$ such that $h(\widetilde{\omega}_{\mathcal{D}}) = \widetilde{\omega}_{\mathcal{C}}$ and $h(\pi_{\mathcal{D}}) = \pi_{\mathcal{C}}$.*

Proof. Let $\mathcal{C} = \langle \widetilde{\omega}_{\mathcal{C}}, \pi_{\mathcal{C}} \rangle$ and $\mathcal{D} = \langle \widetilde{\omega}_{\mathcal{D}}, \pi_{\mathcal{D}} \rangle$ be two CC Lie ideals of \mathcal{L} . Suppose that $g(\widetilde{\omega}_{\mathcal{C}}) = \widetilde{\omega}_{\mathcal{D}}$ and $g(\pi_{\mathcal{C}}) = \pi_{\mathcal{D}}$ for some $g \in \text{Aut}(\mathcal{L})$, then

$$\widetilde{\omega}_{\mathcal{D}}(l_1) = g(\widetilde{\omega}_{\mathcal{C}}) = \sup_{l_2 \in g^{-1}(l_1)} \widetilde{\omega}_{\mathcal{C}}(l_2) = \widetilde{\omega}_{\mathcal{C}}(g^{-1}(l_1))$$

and

$$\pi_{\mathcal{D}}(\Psi(l_1)) = g(\pi_{\mathcal{C}}) = \inf_{l_2 \in g^{-1}(l_1)} \pi_{\mathcal{C}}(l_2) = \pi_{\mathcal{C}}(g^{-1}(l_1)).$$

Thus,

$$g^{-1}(l_1) = \sup_{l_2 \in g(l_1)} \widetilde{\omega}_{\mathcal{D}}(l_2) = \widetilde{\omega}_{\mathcal{D}}(g(l_1)) = \widetilde{\omega}_{\mathcal{C}}(g^{-1}(g(l_1))) = \widetilde{\omega}_{\mathcal{C}}(l_1)$$

and

$$g^{-1}(l_1) = \inf_{l_2 \in g(l_1)} \pi_{\mathcal{D}}(l_2) = \pi_{\mathcal{D}}(g(l_1)) = \pi_{\mathcal{C}}(g^{-1}(g(l_1))) = \pi_{\mathcal{C}}(l_1)$$

for any $l_1 \in \mathcal{L}$. If $h = g^{-1}$, then $h \in \text{Aut}(\mathcal{L})$ and therefore $h(\widetilde{\omega}_{\mathcal{D}}) = \widetilde{\omega}_{\mathcal{C}}$ and $h(\pi_{\mathcal{D}}) = \pi_{\mathcal{C}}$. \square

Theorem 3.19. *If $h(\widetilde{\omega}_{\mathcal{D}}) = \widetilde{\omega}_{\mathcal{C}}$ and $h(\pi_{\mathcal{D}}) = \pi_{\mathcal{C}}$ for some $h \in \text{Aut}(\mathcal{L})$, then $\Upsilon(\widetilde{\omega}_{\mathcal{C}}; \widetilde{\nu}) = h(\Upsilon(\widetilde{\omega}_{\mathcal{D}}; \widetilde{\nu}))$ and $\Upsilon(\pi_{\mathcal{C}}; \gamma) = h(\Upsilon(\pi_{\mathcal{D}}; \gamma))$ for all $\widetilde{\nu} \in \mathbb{P}[0, 1]$ and $\gamma \in [-1, 0]$.*

Proof. Let $\mathcal{C} = \langle \widetilde{\omega}_{\mathcal{C}}, \pi_{\mathcal{C}} \rangle$ and $\mathcal{D} = \langle \widetilde{\omega}_{\mathcal{D}}, \pi_{\mathcal{D}} \rangle$ be a CC Lie ideals of \mathcal{L} . Suppose that $h(\widetilde{\omega}_{\mathcal{D}}) = \widetilde{\omega}_{\mathcal{C}}$ and $h(\pi_{\mathcal{D}}) = \pi_{\mathcal{C}}$ for some $h \in \text{Aut}(\mathcal{L})$, then

$$\widetilde{\omega}_{\mathcal{C}}(l_1) = h(\widetilde{\omega}_{\mathcal{D}})(l_1) = \sup_{l_2 \in h^{-1}(l_1)} \widetilde{\omega}_{\mathcal{D}}(l_2) = \widetilde{\omega}_{\mathcal{D}}(h^{-1}(l_1))$$

and

$$\pi_{\mathcal{C}}(l_1) = h(\pi_{\mathcal{D}})(l_1) = \inf_{l_2 \in h^{-1}(l_1)} \pi_{\mathcal{D}}(l_2) = \pi_{\mathcal{D}}(h^{-1}(l_1))$$

for any $l_1 \in \mathcal{L}$. Now, let $\widetilde{\nu} \in \mathbb{D}[0, 1]$ and $\gamma \in [-1, 0]$. If $l_1 \in \Upsilon(\widetilde{\omega}_{\mathcal{C}}; \widetilde{\nu}) \cap \Upsilon(\pi_{\mathcal{C}}; \gamma)$, then

$$\widetilde{\omega}_{\mathcal{D}}(h^{-1}(l_1)) = \widetilde{\omega}_{\mathcal{C}}(l_1) \geq \widetilde{\nu}$$

and

$$\pi_{\mathcal{D}}(h^{-1}(l_1)) = \pi_{\mathcal{C}}(l_1) \leq \gamma,$$

which implies that $h^{-1}(l_1) \in \Upsilon(\widetilde{\omega}_{\mathcal{D}}; \widetilde{\nu}) \cap \Upsilon(\pi_{\mathcal{D}}; \gamma)$, i.e., $x \in h(\Upsilon(\widetilde{\omega}_{\mathcal{D}}; \widetilde{\nu}) \cap \Upsilon(\pi_{\mathcal{D}}; \gamma))$. Hence, $\Upsilon(\widetilde{\omega}_{\mathcal{C}}; \widetilde{\nu}) \subseteq h(\Upsilon(\widetilde{\omega}_{\mathcal{D}}; \widetilde{\nu}))$ and $\Upsilon(\pi_{\mathcal{C}}; \gamma) \subseteq h(\Upsilon(\pi_{\mathcal{D}}; \gamma))$. Now, let $l_1 \in h(\Upsilon(\widetilde{\omega}_{\mathcal{D}}; \widetilde{\nu})) \cap h(\Upsilon(\pi_{\mathcal{D}}; \gamma))$. Then, $h^{-1}(l_1) \in \Upsilon(\widetilde{\omega}_{\mathcal{D}}; \widetilde{\nu}) \cap \Upsilon(\pi_{\mathcal{D}}; \gamma)$. Therefore,

$$\widetilde{\omega}_{\mathcal{C}}(l_1) = \widetilde{\omega}_{\mathcal{D}}(h^{-1}(l_1)) \geq \widetilde{\nu}$$

and

$$\pi_{\mathcal{C}}(l_1) = \pi_{\mathcal{D}}(h^{-1}(l_1)) \leq \gamma.$$

It follows that $l_1 \in \Upsilon(\widetilde{\omega}_{\mathcal{C}}; \widetilde{\nu}) \cap \Upsilon(\pi_{\mathcal{C}}; \gamma)$. Hence, $h(\Upsilon(\widetilde{\omega}_{\mathcal{D}}; \widetilde{\nu})) \subseteq \Upsilon(\widetilde{\omega}_{\mathcal{C}}; \widetilde{\nu})$ and $h(\Upsilon(\pi_{\mathcal{D}}; \gamma)) \subseteq \Upsilon(\pi_{\mathcal{C}}; \gamma)$. Thus, $\Upsilon(\widetilde{\omega}_{\mathcal{C}}; \widetilde{\nu}) = h(\Upsilon(\widetilde{\omega}_{\mathcal{D}}; \widetilde{\nu}))$ and $\Upsilon(\pi_{\mathcal{C}}; \gamma) = h(\Upsilon(\pi_{\mathcal{D}}; \gamma))$ for all $\widetilde{\nu} \in \mathbb{P}[0, 1]$ and $\gamma \in [-1, 0]$. \square

Theorem 3.20. *If $\exists h \in \text{Aut } \mathcal{L}$ such that $\Upsilon(\widetilde{\omega}_{\mathcal{C}}; \widetilde{\nu}) = h(\Upsilon(\widetilde{\omega}_{\mathcal{D}}; \widetilde{\nu}))$ and $\Upsilon(\pi_{\mathcal{C}}; \gamma) = h(\Upsilon(\pi_{\mathcal{D}}; \gamma))$ for all $(\widetilde{\nu}, \gamma) \in \mathbb{P}[0, 1] \times [-1, 0]$, then \mathcal{C} is a crossing structure having the same type of \mathcal{D} .*

Proof. Let $\mathcal{C} = \langle \widetilde{\omega}_{\mathcal{C}}, \pi_{\mathcal{C}} \rangle$ and $\mathcal{D} = \langle \widetilde{\omega}_{\mathcal{D}}, \pi_{\mathcal{D}} \rangle$ be two CC Lie ideals of \mathcal{L} . Assume that $\exists h \in \text{Aut } \mathcal{L}$ such that $\Upsilon(\widetilde{\omega}_{\mathcal{C}}; \widetilde{\nu}) = h(\Upsilon(\widetilde{\omega}_{\mathcal{D}}; \widetilde{\nu}))$ and $\Upsilon(\pi_{\mathcal{C}}; \gamma) = h(\Upsilon(\pi_{\mathcal{D}}; \gamma))$ for all $(\widetilde{\nu}, \gamma) \in \mathbb{P}[0, 1] \times [-1, 0]$. Let $\widetilde{\omega}_{\mathcal{D}}(\mathcal{L}^{-1}(l_1)) = \widetilde{\nu}$ and $\omega_{\mathcal{D}}(\mathcal{L}^{-1}(l_1)) = \gamma$. Then, $h^{-1}(l_1) \in \Upsilon(\widetilde{\omega}_{\mathcal{C}}; \widetilde{\nu}) \cap \Upsilon(\pi_{\mathcal{C}}; \gamma)$. Hence, $l_1 \in h(\Upsilon(\widetilde{\omega}_{\mathcal{D}}; \widetilde{\nu})) \cap h(\Upsilon(\pi_{\mathcal{D}}; \gamma)) = \Upsilon(\widetilde{\omega}_{\mathcal{C}}; \widetilde{\nu}) \cap \Upsilon(\pi_{\mathcal{C}}; \gamma)$. Thus, $\widetilde{\omega}_{\mathcal{C}}(l_1) \geq \widetilde{\nu} = \widetilde{\omega}_{\mathcal{D}}(h^{-1}(l_1))$ and $\omega_{\mathcal{C}}(l_1) \leq \gamma = \omega_{\mathcal{D}}(h^{-1}(l_1))$ for all $l_1 \in \mathcal{L}$. Therefore, \mathcal{C} is a crossing structure having the same type of \mathcal{D} . \square

4. Crossing cubic quotient Lie algebras

Here, we give a construction of a quotient Lie algebra via CC Lie ideal \mathcal{L} . Then, we present the CC isomorphism theorems.

Theorem 4.1. *Let \mathcal{I} be a Lie ideal of \mathcal{L} . If $\mathcal{C} = \langle \widetilde{\omega}_{\mathcal{C}}, \pi_{\mathcal{C}} \rangle$ is a CC Lie ideal of \mathcal{L} , then a CCS $\overline{\mathcal{C}} = \langle \overline{\widetilde{\omega}_{\mathcal{C}}}, \overline{\pi_{\mathcal{C}}} \rangle$ is a CC Lie ideal of the quotient Lie algebra \mathcal{L}/\mathcal{I} , where $\overline{\widetilde{\omega}_{\mathcal{C}}} = \sup_{k \in \mathcal{I}}(l_1 + k)$ and $\overline{\pi_{\mathcal{C}}} = \inf_{k \in \mathcal{I}}(l_1 + k)$.*

Proof. Straightforward, $\overline{\mathcal{C}}$ is well defined. Let $(l_1 + \mathcal{I}), (l_2 + \mathcal{I}) \in \mathcal{L}/\mathcal{I}$, then

$$\begin{aligned} \overline{\widetilde{\omega}_{\mathcal{C}}}((l_1 + \mathcal{I}) + (l_2 + \mathcal{I})) &= \overline{\widetilde{\omega}_{\mathcal{C}}}((l_1 + l_2) + \mathcal{I}) \\ &= \sup_{k \in \mathcal{I}}((l_1 + l_2) + k) \\ &= \sup_{k = \alpha + \beta \in \mathcal{I}}((l_1 + l_2) + k) \\ &\geq \sup_{\alpha, \beta \in \mathcal{I}} \min\{\overline{\widetilde{\omega}_{\mathcal{C}}}(l_1 + \alpha), \overline{\widetilde{\omega}_{\mathcal{C}}}(l_2 + \beta)\} \\ &= \min_{\alpha \in \mathcal{I}} \sup(l_1 + \alpha), \min_{\beta \in \mathcal{I}} \sup(l_2 + \beta) \\ &= \min\{\overline{\widetilde{\omega}_{\mathcal{C}}}(l_1 + \mathcal{I}), \overline{\widetilde{\omega}_{\mathcal{C}}}(l_2 + \mathcal{I})\}, \end{aligned}$$

$$\overline{\widetilde{\omega}_{\mathcal{C}}}(\delta(l_1 + \mathcal{I})) = \overline{\widetilde{\omega}_{\mathcal{C}}}(\delta l_1 + \mathcal{I}) = \sup_{k \in \mathcal{I}}(\delta l_1 + k) \geq \sup_{k \in \mathcal{I}}(l_1 + k) = \overline{\widetilde{\omega}_{\mathcal{C}}}(l_1 + \mathcal{I})$$

and

$$\begin{aligned}
 \overline{\widetilde{\omega}_{\mathcal{C}}}([l_1 + \mathcal{I}], [l_2 + \mathcal{I}]) &= \overline{\widetilde{\omega}_{\mathcal{C}}}([l_1 + l_2] + \mathcal{I}) \\
 &= \sup_{k \in \mathcal{I}} \overline{\widetilde{\omega}_{\mathcal{C}}}([l_1 + l_2] + k) \\
 &\geq \sup_{k \in \mathcal{I}} \overline{\widetilde{\omega}_{\mathcal{C}}}(l_1 + k) \\
 &= \overline{\widetilde{\omega}_{\mathcal{C}}}(l_1 + \mathcal{I}).
 \end{aligned}$$

Also,

$$\begin{aligned}
 \overline{\pi_{\mathcal{C}}}((l_1 + \mathcal{I}) + (l_2 + \mathcal{I})) &= \overline{\pi_{\mathcal{C}}}((l_1 + l_2) + \mathcal{I}) \\
 &= \inf_{k \in \mathcal{I}} ((l_1 + l_2) + k) \\
 &= \inf_{k = \alpha + \beta \in \mathcal{I}} ((l_1 + l_2) + k) \\
 &\leq \inf_{\alpha, \beta \in \mathcal{I}} \max\{\overline{\pi_{\mathcal{C}}}(l_1 + \alpha), \overline{\pi_{\mathcal{C}}}(l_2 + \beta)\} \\
 &= \max\{\inf_{\alpha \in \mathcal{I}} (l_1 + \alpha), \inf_{\beta \in \mathcal{I}} (l_2 + \beta)\} \\
 &= \max\{\overline{\pi_{\mathcal{C}}}(l_1 + \mathcal{I}), \overline{\pi_{\mathcal{C}}}(l_2 + \mathcal{I})\},
 \end{aligned}$$

$$\overline{\pi_{\mathcal{C}}}(\delta(l_1 + \mathcal{I})) = \overline{\pi_{\mathcal{C}}}(\delta l_1 + \mathcal{I}) = \inf_{k \in \mathcal{I}} (\delta l_1 + k) \leq \inf_{k \in \mathcal{I}} (l_1 + k) = \overline{\pi_{\mathcal{C}}}(l_1 + \mathcal{I})$$

and

$$\begin{aligned}
 \overline{\pi_{\mathcal{C}}}([l_1 + \mathcal{I}], [l_2 + \mathcal{I}]) &= \overline{\pi_{\mathcal{C}}}([l_1 + l_2] + \mathcal{I}) \\
 &= \inf_{k \in \mathcal{I}} \overline{\pi_{\mathcal{C}}}([l_1 + l_2] + k) \\
 &\leq \inf_{k \in \mathcal{I}} \overline{\pi_{\mathcal{C}}}(l_1 + k) \\
 &= \overline{\pi_{\mathcal{C}}}(l_1 + \mathcal{I}).
 \end{aligned}$$

Hence, $\overline{\mathcal{C}}$ is a CC Lie ideal of \mathcal{L}/\mathcal{I} . □

Theorem 4.2. Let $\Psi : \mathcal{L} \rightarrow \mathcal{L}$ be an epimorphism of a Lie algebra \mathcal{L} onto a Lie algebra \mathcal{L} . Then, the following statements hold:

- (i) If \mathcal{C} is a CC Lie ideal of \mathcal{L} , then $\Psi(\mathcal{C})$ is a CC Lie ideal of \mathcal{L} .
- (ii) If \mathcal{D} is a CC Lie ideal of \mathcal{L} , then $\Psi^{-1}(\mathcal{D})$ is a CC Lie ideal of \mathcal{L} .

Proof. Straightforward. □

An equivalence relation \bowtie in Theorem 3.5 is a congruence relation. To verify that \bowtie is a congruence relation on \mathcal{L} . Let $l_1 \bowtie l_2$ and $l_2 \bowtie l_3$, then $\widetilde{\omega}_{\mathcal{C}}(l_1 - l_2) = \widetilde{\omega}_{\mathcal{C}}(0)$, $\widetilde{\omega}_{\mathcal{C}}(l_2 - l_3) = \widetilde{\omega}_{\mathcal{C}}(0)$, $\pi_{\mathcal{C}}(l_1 - l_2) = \pi_{\mathcal{C}}(0)$ and $\pi_{\mathcal{C}}(l_2 - l_3) = \pi_{\mathcal{C}}(0)$. Now, let $n_1, n_2, m_1, m_2 \in \mathcal{L}$, we have

$$\begin{aligned}
 \widetilde{\omega}_{\mathcal{C}}((n_1 + n_2) - (m_1 + m_2)) &= \widetilde{\omega}_{\mathcal{C}}((n_1 - m_1) + (n_2 - m_2)) \\
 &\geq \widetilde{\min}\{\widetilde{\omega}_{\mathcal{C}}(n_1 - m_1), \widetilde{\omega}_{\mathcal{C}}(n_2 - m_2)\}
 \end{aligned}$$

$$= \widetilde{\omega}_{\mathcal{C}}(0),$$

$$\widetilde{\omega}_{\mathcal{C}}(\alpha n_1 - \alpha m_1) = \widetilde{\omega}_{\mathcal{C}}(\alpha(n_1 - m_1)) \geq \widetilde{\omega}_{\mathcal{C}}(n_1 - m_1) = \widetilde{\omega}_{\mathcal{C}}(0),$$

$$\begin{aligned} \widetilde{\omega}_{\mathcal{C}}([n_1, n_2] - [m_1, m_2]) &= \widetilde{\omega}_{\mathcal{C}}([n_1 - m_1], [n_2 - m_2]) \\ &\geq \widetilde{\min}\{\widetilde{\omega}_{\mathcal{C}}(n_1 - m_1), \widetilde{\omega}_{\mathcal{C}}(n_2 - m_2)\} \\ &= \widetilde{\omega}_{\mathcal{C}}(0). \end{aligned}$$

Also,

$$\begin{aligned} \pi_{\mathcal{C}}((n_1 + n_2) - (m_1 + m_2)) &= \pi_{\mathcal{C}}((n_1 - m_1) + (n_2 - m_2)) \\ &\leq \max\{\pi_{\mathcal{C}}(n_1 - m_1), \pi_{\mathcal{C}}(n_2 - m_2)\} \\ &= \pi_{\mathcal{C}}(0), \end{aligned}$$

$$\pi_{\mathcal{C}}(\alpha n_1 - \alpha m_1) = \pi_{\mathcal{C}}(\alpha(n_1 - m_1)) \leq \pi_{\mathcal{C}}(n_1 - m_1) = \pi_{\mathcal{C}}(0),$$

$$\begin{aligned} \pi_{\mathcal{C}}([n_1, n_2] - [m_1, m_2]) &= \pi_{\mathcal{C}}([n_1 - m_1], [n_2 - m_2]) \\ &\leq \max\{\pi_{\mathcal{C}}(n_1 - m_1), \pi_{\mathcal{C}}(n_2 - m_2)\} \\ &= \pi_{\mathcal{C}}(0). \end{aligned}$$

That is, $n_1 + n_2 \bowtie m_1 + m_2$, $\alpha n_1 \bowtie \alpha m_1$ and $[n_1, n_2] \bowtie [m_1, m_2]$. Thus, \bowtie is a congruence relation on \mathcal{L} .

The set of all equivalence classes $\mathcal{C}[\mathcal{L}]$ is denoted by \mathcal{L}/\mathcal{C} . It is a Lie algebra under the following operations:

$$\begin{aligned} \overline{\widetilde{\omega}_{\mathcal{C}}}[\mathcal{L}] + \overline{\widetilde{\omega}_{\mathcal{C}}}[\mathcal{L}] &= \overline{\widetilde{\omega}_{\mathcal{C}}}[\mathcal{L} + \mathcal{L}], \alpha \overline{\widetilde{\omega}_{\mathcal{C}}}[\mathcal{L}] = \overline{\widetilde{\omega}_{\mathcal{C}}}[\alpha \mathcal{L}], [\overline{\widetilde{\omega}_{\mathcal{C}}}[\mathcal{L}], \overline{\widetilde{\omega}_{\mathcal{C}}}[\mathcal{L}]] = \overline{\widetilde{\omega}_{\mathcal{C}}}[[\mathcal{L} + \mathcal{L}]], \\ \overline{\pi_{\mathcal{C}}}[\mathcal{L}] + \overline{\pi_{\mathcal{C}}}[\mathcal{L}] &= \overline{\pi_{\mathcal{C}}}[\mathcal{L} + \mathcal{L}], \alpha \overline{\pi_{\mathcal{C}}}[\mathcal{L}] = \overline{\pi_{\mathcal{C}}}[\alpha \mathcal{L}] \text{ and } [\overline{\pi_{\mathcal{C}}}[\mathcal{L}], \overline{\pi_{\mathcal{C}}}[\mathcal{L}]] = \overline{\pi_{\mathcal{C}}}[[\mathcal{L} + \mathcal{L}]]. \end{aligned}$$

Next, we present the CC isomorphism theorems over Lie algebras.

Theorem 4.3. Let $\Psi : \mathcal{L} \rightarrow \mathcal{L}'$ be an epimorphism, where \mathcal{L} and \mathcal{L}' are Lie algebras. If $\mathcal{C} = \langle \widetilde{\omega}_{\mathcal{C}}, \pi_{\mathcal{C}} \rangle$ is a CC Lie ideal in \mathcal{L} , then $\mathcal{L}/\Psi^{-1}(\mathcal{C}) \cong \mathcal{L}'/\mathcal{C}$.

Proof. Define $\Omega : \mathcal{L}/\Psi^{-1}(\mathcal{C}) \rightarrow \mathcal{L}'/\mathcal{C}$ as:

$$\Omega(\Psi^{-1}(\widetilde{\omega}_{\mathcal{C}}(l))) = \widetilde{\omega}_{\mathcal{C}}(\Psi(l)) \text{ and } \Omega(\Psi^{-1}(\pi_{\mathcal{C}}(l))) = \pi_{\mathcal{C}}(\Psi(l)).$$

$\forall l \in \mathcal{L}$ and for any collection of the CCSs of \mathcal{L}' . Hence, Ω is an isomorphism. \square

Theorem 4.4. Let $\mathcal{C} = \langle \widetilde{\omega}_{\mathcal{C}}, \pi_{\mathcal{C}} \rangle$ and $\mathcal{D} = \langle \widetilde{\omega}_{\mathcal{D}}, \pi_{\mathcal{D}} \rangle$ be two CC Lie ideals in \mathcal{L} , with $\widetilde{\omega}_{\mathcal{C}}(0) = \widetilde{\omega}_{\mathcal{D}}(0)$ and $\pi_{\mathcal{C}}(0) = \pi_{\mathcal{D}}(0)$. Then, $(\mathcal{L}_{\mathcal{C}} + \mathcal{L}_{\mathcal{D}})/\mathcal{D} \cong \mathcal{L}_{\mathcal{C}}/(\mathcal{C} \cap \mathcal{D})$.

Proof. Clearly, $\mathcal{L}_{\mathcal{C}} + \mathcal{L}_{\mathcal{D}}$ and $\mathcal{L}_{\mathcal{C}}/(\mathcal{C} \cap \mathcal{D})$ are CC Lie ideals in \mathcal{L} . For every $l \in (\mathcal{L}_{\mathcal{C}} + \mathcal{L}_{\mathcal{D}})$, $l = a + b$, where $a \in \mathcal{L}_{\mathcal{C}}$ and $b \in \mathcal{L}_{\mathcal{D}}$. Let \mathcal{C} and \mathcal{D} have a congruence relation \propto on \mathcal{L} , defined as:

$$a \propto w \Leftrightarrow (\widetilde{\omega}_{\mathcal{C}}(a - w) = \widetilde{\omega}_{\mathcal{C}}(0), \pi_{\mathcal{C}}(a - w) = \pi_{\mathcal{C}}(0), \widetilde{\omega}_{\mathcal{D}}(a - w) = \widetilde{\omega}_{\mathcal{D}}(0), \pi_{\mathcal{D}}(a - w) = \pi_{\mathcal{D}}(0))$$

for all \mathcal{C} and \mathcal{D} . Define $\Omega : (\mathcal{L}_{\mathcal{C}} + \mathcal{L}_{\mathcal{D}})/\mathcal{D} \rightarrow \mathcal{L}_{\mathcal{C}}/(\mathcal{C} \cap \mathcal{D})$ as:

$$\Omega((\widetilde{\omega}_{\mathcal{D}}(l))) = \Omega(\widetilde{\omega}_{\mathcal{D}}(a + b)) = (\widetilde{\omega}_{\mathcal{C}} \cap \widetilde{\omega}_{\mathcal{D}})(a)$$

and

$$\Omega((\pi_{\mathcal{D}}(l))) = \Omega(\pi_{\mathcal{D}}(a + b)) = (\pi_{\mathcal{C}} \cap \pi_{\mathcal{D}})(a).$$

Then, Ω is an isomorphism. \square

Theorem 4.5. Let $\mathcal{C} = \langle \widetilde{\omega}_{\mathcal{C}}, \pi_{\mathcal{C}} \rangle$ and $\mathcal{D} = \langle \widetilde{\omega}_{\mathcal{D}}, \pi_{\mathcal{D}} \rangle$ be two CC Lie ideals in \mathcal{L} , with $\widetilde{\omega}_{\mathcal{C}}(0) = \widetilde{\omega}_{\mathcal{D}}(0)$, $\pi_{\mathcal{C}}(0) = \pi_{\mathcal{D}}(0)$ and $\mathcal{C} \subseteq \mathcal{D}$. Then, $(\mathcal{L}/\mathcal{C})/(\mathcal{L}_{\mathcal{D}}/\mathcal{C}) \cong \mathcal{L}/(\mathcal{D})$.

Proof. Clearly, $(\mathcal{L}/\mathcal{C})/(\mathcal{L}_{\mathcal{D}}/\mathcal{C})$ is a CC Lie ideal over \mathcal{L} . Define

$$\Omega : (\mathcal{L}/\mathcal{C})/(\mathcal{L}_{\mathcal{D}}/\mathcal{C}) \rightarrow \mathcal{L}/(\mathcal{D})$$

as:

$$\Omega((\widetilde{\omega}_{\mathcal{C}}(l) + (\mathcal{L}_{\mathcal{D}}/\mathcal{C}))) = \widetilde{\omega}_{\mathcal{D}}(l) \text{ and } \Omega((\pi_{\mathcal{C}}(l) + (\mathcal{L}_{\mathcal{D}}/\mathcal{C}))) = \pi_{\mathcal{D}}(l)$$

$\forall l \in \mathcal{L}$. Then, the proof is straightforward. \square

5. Conclusions

As discussed earlier, the CC structure is a parallel concept between interval-valued and negative versions of fuzziness structures. The idea of the CC structure is an extension of the fuzziness structure of two polarities (positive and negative). In this paper, we develop the concepts of CC Lie subalgebras, CC Lie ideals, and homomorphism. We discussed the Cartesian product of CC Lie subalgebras. Also, we described some interesting results of CC Lie ideals. In addition, we provided a construction of a quotient Lie algebra via the CC Lie ideal in a Lie algebra and presented the CC isomorphism theorems.

The idea of CC structure is a topic of great interest, and it is a task to apply this logical concept to problems in theoretical physics, coding theory, and quantum mechanics. Further, the results of this study can be utilized in a variety of algebraic frameworks, for instance Hom-Lie algebras, Hom-Lie bi-algebras, semirings, Γ -semirings, and QS/UP/KU-algebras.

Author contributions

Anas Al-Masarwah: Conceptualization and Methodology. Nadeen Kdaisat and Majdoleen Abuqamar: Investigation and Writing-original draft. Anas Al-Masarwah and Kholood Alsager: Writing-review and editing. All authors have read and agreed to the published version of the manuscript.

Acknowledgments

The Researchers would like to thank the Deanship of Graduate Studies and Scientific Research at Qassim University for financial support (QU-APC-2024-9/1).

Conflict of interest

The authors declare that they have no competing interests.

References

1. L. Dorst, C. Doran, J. Lasenby, *Applications of geometric algebra in computer science and engineering*, Springer Science & Business Media: Berlin/Heidelberg, Germany, 2012.
2. W. Wechler, *Universal algebra for computer scientists*, Springer Science & Business Media: Berlin/Heidelberg, Germany, 2012.
3. L. Dorst, D. Fontijne, S. Mann, *Geometric algebra for computer science: an object-oriented approach to geometry*, Elsevier: Amsterdam, The Netherlands, 2007.
4. J. A. de Azcárraga, J. M. Izquierdo, *Lie groups, Lie algebras, cohomology and some applications in physics*, Cambridge University Press: Cambridge, UK, 1998.
5. M. Guerreiro, Group algebras and coding theory, *São Paulo J. Math. Sci.*, **10** (2016), 346–371. <https://doi.org/10.1007/s40863-016-0040-x>
6. J. E. Humphreys, *Introduction to Lie algebras and representation theory*, Vol. 9, Graduate Texts in Mathematics, New York: Springer-Verlag, 1972. <https://doi.org/10.1007/978-1-4612-6398-2>
7. L. A. Zadeh, Fuzzy sets, *Inform. Control*, **8** (1965), 338–353. [https://doi.org/10.1016/S0019-9958\(65\)90241-X](https://doi.org/10.1016/S0019-9958(65)90241-X)
8. L. A. Zadeh, The concept of a linguistic variable and its application to approximate reasoning–I, *Inform. Sciences*, **8** (1975), 199–249. [https://doi.org/10.1016/0020-0255\(75\)90036-5](https://doi.org/10.1016/0020-0255(75)90036-5)
9. Y. B. Jun, C. S. Kim, K. O. Yang, Cubic sets, *Ann. Fuzzy Math. Inform.*, **4** (2012), 83–98.
10. Y. B. Jun, K. J. Lee, S. Z. Song, N -ideals of BCK/BCI-algebras, *J. Chungcheong Math. Soc.*, **22** (2009), 417–437.
11. A. Fallatah, M. O. Massa'deh, A. U. Alkouri, Normal and cosets of (γ, δ) -fuzzy HX-subgroups, *J. Appl. Math. Inform.*, **40** (2022), 719–727. <https://doi.org/10.14317/jami.2022.719>
12. A. Al-Masarwah, A. G. Ahmad, m -polar fuzzy ideals of BCK/BCI-algebras, *J. King Saud Univ. Sci.*, **31** (2019), 1220–1226. <https://doi.org/10.1016/j.jksus.2018.10.002>
13. E. A. Abuhijleh, M. Massa'deh, A. Sheimat, A. Alkouri, Complex fuzzy groups based on Rosenfeld's approach, *WSEAS Trans. Math.*, **20** (2021), 368–377. <https://doi.org/10.37394/23206.2021.20.38>
14. G. Muhiuddin, M. E. Elnair, D. Al-Kadi, N -structures applied to commutative ideals of BCI-algebras, *Symmetry*, **14** (2022), 2015. <https://doi.org/10.3390/sym14102015>
15. G. Muhiuddin, K. Porselvi, B. Elavarasan, D. Al-Kadi, Neutrosophic K -structures in ordered semigroups, *Comput. Model. Eng. Sci.*, **131** (2022), 979–999. <https://doi.org/10.32604/cmesci.2022.018615>
16. A. Mahboob, G. Muhiuddin, m -polar cubic $p(q$ and $a)$ -ideals of BCI-algebras, *Discrete Math., Algorithms Appl.*, **16** (2024), 2350013. <https://doi.org/10.1142/S1793830923500131>

17. G. Muhiuddin, D. Al-Kadi, A. Mahboob, Hybrid structures applied to ideals in BCI-algebras, *J. Math.*, **2020** (2020), 1–7. <https://doi.org/10.1155/2020/2365078>
18. S. E. Yehia, Fuzzy ideals and fuzzy subalgebras of Lie algebras, *Fuzzy Sets Syst.*, **80** (1996), 237–244. [https://doi.org/10.1016/0165-0114\(95\)00109-3](https://doi.org/10.1016/0165-0114(95)00109-3)
19. M. Akram, Generalized fuzzy Lie subalgebras, In: *Fuzzy Lie algebras*, Singapore: Springer, 2018. https://doi.org/10.1007/978-981-13-3221-0_4
20. M. Akram, K. P. Shum, Fuzzy Lie ideals over a fuzzy field, *Ital. J. Pure Appl. Math.*, **27** (2010), 281–292.
21. M. Akram, *Fuzzy Lie algebra*, Springer Nature, 2018. <https://doi.org/10.1007/978-981-13-3221-0>
22. M. Akram, Fuzzy Lie ideals of Lie algebras with interval-valued membership functions, *Quasigroups Related Syst.*, **16** (2008), 1–12.
23. M. Akram, K. P. Shum, Intuitionistic fuzzy Lie algebras, *Southeast Asian Bull. Math.*, **31** (2005), 843–855.
24. M. Akram, H. Gulzar, K. P. Shum, Single-valued neutrosophic Lie algebras, *J. Math. Res. Appl.*, **39** (2019), 141–152. <https://doi.org/10.3770/j.issn:2095-2651.2019.02.003>
25. Y. B. Jun, K. Hur, J. G. Lee, J. Kim, Crossing cubic structures as an extension of bipolar fuzzy sets, *Ann. Fuzzy Math. Inform.*, **22** (2021), 1–15. <https://doi.org/10.30948/afmi.2021.22.1.1>
26. Y. B. Jun, S. Z. Song, Crossing cubic ideals of BCK/BCI-algebras, *J. Algebraic Hyperstructures Logical Algebras*, **2** (2021), 17–31. <https://doi.org/10.52547/HATEF.JAHLA.2.1.2>
27. M. A. Öztürk, D. Yılmaz, Y. B. Jun, Semigroup structures and commutative ideals of BCK-algebras based on crossing cubic set structures, *Axioms*, **11** (2022), 25. <https://doi.org/10.3390/axioms11010025>
28. S. M. Mostafa, J. G. Lee, Y. B. Jun, K. Hur, C. K. S. Ebid, $(\widetilde{\alpha}, \alpha)$ -crossing cubic QS-ideal of QS-algebra, *Ann. Fuzzy Math. Inform.*, **24** (2022), 55–73. <https://doi.org/10.30948/afmi.2022.24.1.55>



AIMS Press

© 2024 the Author(s), licensee AIMS Press. This is an open access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>)